

PUBLISHED VERSION

Rajesh Chaudhary, Atip Likidlilid, Thavatchai Peerapatdit, Damras Tresukosol, Sorachai Srisuma, Suphachai Ratanamaneechat and Charn Sriratanasathavorn

Apolipoprotein E gene polymorphism: effects on plasma lipids and risk of type 2 diabetes and coronary artery disease

Cardiovascular Diabetology, 2012; 11(1):36-1-36-11

© 2012 Chaudhary et al; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Originally published at <http://dx.doi.org/10.1186/1475-2840-11-36>

PERMISSIONS

<https://creativecommons.org/licenses/by/2.0/>



The image shows the Creative Commons Attribution 2.0 Generic (CC BY 2.0) license logo and text. The logo consists of two circles: one with 'cc' and another with a person icon. Below the logo, the text reads 'Attribution 2.0 Generic (CC BY 2.0)'. A disclaimer states: 'This is a human-readable summary of (and not a substitute for) the license. [Disclaimer.](#)'. The main text is divided into two sections: 'You are free to:' and 'Under the following terms:'. Under 'You are free to:', there are two bullet points: 'Share — copy and redistribute the material in any medium or format' and 'Adapt — remix, transform, and build upon the material for any purpose, even commercially.' A circular seal on the right says 'Free Cultural APPROVED FOR Works'. Below this, it states: 'The licensor cannot revoke these freedoms as long as you follow the license terms.' Under 'Under the following terms:', there are two bullet points: 'Attribution — You must give [appropriate credit](#), provide a link to the license, and [indicate if changes were made](#). You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.' and 'No additional restrictions — You may not apply legal terms or [technological measures](#) that legally restrict others from doing anything the license permits.'

20 June 2017

<http://hdl.handle.net/2440/106095>

ORIGINAL INVESTIGATION

Open Access

Apolipoprotein E gene polymorphism: effects on plasma lipids and risk of type 2 diabetes and coronary artery disease

Rajesh Chaudhary¹, Atip Likidilid^{1*}, Thavatchai Peerapatdit², Damras Tresukosol², Sorachai Srisuma³, Suphachai Ratanamaneechat⁴ and Charn Sriratanasathavorn⁵

Abstract

Background: The most common apolipoprotein E (apoE) gene polymorphism has been found to influence plasma lipid concentration and its correlation with coronary artery disease (CAD) has been extensively investigated in the last decade. It is, however, unclear whether apoE gene polymorphism is also associated with increased risk of type 2 diabetes mellitus (T2DM). The knowledge of this study may provide the primary prevention for T2DM and CAD development before its initiation and progression. Therefore, this study was carried out to determine the association between apoE gene polymorphism and T2DM with and without CAD and its role in lipid metabolism.

Methods: The case-control study was carried out on a total of 451 samples including 149 normal control subjects, 155 subjects with T2DM, and 147 subjects with T2DM complicated with CAD. The apoE gene polymorphism was tested by polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP). Univariable and multivariable logistic regression analyses were used to identify the possible risks of T2DM and CAD.

Results: A significantly increased frequency of E3/E4 genotype was observed only in T2DM with CAD group ($p = 0.0004$), whereas the $\epsilon 4$ allele was significantly higher in both T2DM ($p = 0.047$) and T2DM with CAD ($p = 0.009$) as compared with controls. E3/E4 genotype was also the independent risk in developing CAD after adjusting with established risk factors with adjusted odds ratio (OR) 2.52 (95%CI 1.28-4.97, $p = 0.008$). The independent predictor of individuals carrying $\epsilon 4$ allele still remained significantly associated with both CAD (adjusted OR 2.32, 95%CI 1.17-4.61, $p = 0.016$) and T2DM (adjusted OR 2.04, 95%CI 1.07-3.86, $p = 0.029$). After simultaneously examining the joint association of E3/E4 genotype combined with either obesity or smoking the risk increased to approximately 5-fold in T2DM (adjusted OR 4.93, 95%CI 1.74-13.98, $p = 0.003$) and 10-fold in CAD (adjusted OR 10.48, 95%CI 3.56-30.79, $p < 0.0001$). The association between apoE genotypes on plasma lipid levels was compared between E3/E3 as a reference and E4-bearing genotypes. E4-bearing genotypes showed lower HDL-C and higher VLDL-C and TG, whereas other values of plasma lipid concentrations showed no significant difference.

Conclusions: These results indicate that $\epsilon 4$ allele has influence on lipid profiles and is associated with the development of both T2DM with and without CAD, and furthermore, it increased the risk among the subjects with obesity and/or smoking, the conditions associated with high oxidative stress.

Keywords: Apolipoprotein E, Polymorphism, Type 2 diabetes mellitus, Hyperglycemia, Coronary artery disease, Restriction fragment length polymorphism

* Correspondence: atip.lik@mahidol.ac.th

¹Department of Biochemistry, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok 10700, Thailand

Full list of author information is available at the end of the article

Introduction

Type 2 diabetes mellitus (T2DM) is one of the most common diseases with a high incidence and prevalence throughout the world. It affects nearly 4% of the world's population and this percentage will supposedly be increasing up to 5.4% by year 2025 [1]. Prevalence of diabetes in Thai adults as shown by the previous study on Thai population was 9.6% (2.4 million population) and the impaired fasting glucose was 5.4% (1.4 million people). Mean fasting plasma glucose level by age, sex, and area of residence was found to be substantially higher in urban population group than rural [2]. T2DM is also known as a major independent risk factor for coronary artery disease (CAD) and is the major cause of morbidity and mortality affecting people with diabetes. To date, several mechanisms such as dyslipoproteinemia, obesity, oxidative stress, smoking, exercise, alcohol intake, and genetic factors have been identified as risk factors of both T2DM and CAD. Lack of apolipoprotein E (apoE) gene has been clearly demonstrated as a leading cause of severe hyperlipidemia and spontaneous development of atherosclerosis in mammals [3,4]. However, few studies have been able to demonstrate an association between T2DM and various single nucleotide polymorphisms (SNPs) [5]. Recently, Zeljko et al. 2011 [6] indicated that apoE gene polymorphism is also associated with obesity in normal Croatian Roma population. Adipocytes in an obese person which are the central and causal components in T2DM can generate high amount of biologically active molecules called adipokines or adipocytokines such as plasminogen activator inhibitor-1 (PAI-1), resistin, leptin, interleukin-6 (IL-6), and tumor necrosis factor alpha (TNF- α) [7]. These inflammatory cytokines inhibit insulin-stimulated glucose metabolism in skeletal muscles and stimulate gluconeogenesis in hepatocytes causing hyperglycemia [8]. Hyperglycemia-induced oxidative stress results in reducing glucose uptake from blood by muscle cells and develops into insulin resistance by decreasing insulin secretion from pancreatic β -cells [9]. Taken these together, increased oxidative stress in hyperglycemia and reduced lipid clearance because of apoE gene polymorphism are effective in developing insulin resistance and T2DM. In addition, high oxidative stress can also cause vascular inflammation leading to atherosclerosis through several cytokines such as NF- κ B, TNF- α , IL-1 β , and other proinflammatory cytokines [10]. All of these factors are one of the underlying causes of metabolic syndrome [7]. Grundy reported that subjects suffering from the metabolic syndrome are at 2-fold higher risk of developing CAD and at 5-fold higher risk of developing T2DM; however, persons suffering from T2DM are at 3-fold higher risk of developing CAD [11].

ApoE gene is one of the most studied genes which is responsible for stabilizing and solubilizing circulating

lipoproteins in our body and also responsible for the development of CAD [10,11]. ApoE acts as a high affinity ligand for several hepatic lipoprotein receptors such as low-density lipoprotein receptor (LDLR) and LDL-related protein (LRP) and is involved in the process of cellular incorporation of several lipoproteins for transport and digestion [12]. ApoE is a plasma glycoprotein of 34 kDa with 299-amino acids associated with several other plasma glycoproteins, such as high density lipoprotein (HDL), very low density lipoprotein (VLDL), and chylomicrons [13]. In humans, apoE gene located on the chromosome at position 19q13.2 has been known to be polymorphic. SNPs at positions 112 (rs 429358) and 158 (rs 7412) determine three major alleles: ϵ 2 (T to C substitution at position 158), the most common ϵ 3, and ϵ 4 (C to T substitution at position 112); 3 isoforms: ApoE2 (Cys112, 158Cys), ApoE3 (Cys112, 158Arg), and ApoE4 (Arg112, 158Arg); and 6 genotypes having 3 homozygous: E2/E2, E3/E3, E4/E4, and 3 heterozygous: E2/E3, E2/E4, E3/E4 [13]. Previous studies have shown that apoE alleles have influence on the lipid clearance and metabolism in humans. ApoE ϵ 2 allele has been reported to be associated with higher plasma levels of apoE, decreased plasma levels of LDL cholesterol (LDL-C) and lower risk of CAD [14] while apoE ϵ 4 is associated with lower plasma level of apoE, increased plasma levels of total cholesterol (TC), LDL-C, VLDL cholesterol (VLDL-C), and greater risk of CAD when compared to apoE3 homozygotes [15]. One reason for this impaired clearance by apoE ϵ 4 leading to pathogenesis of CAD might be that apoE ϵ 4 binds strongly to LDLR compared to other genotypes. The resulting high amount of lipid can suppress the synthesis of LDLR leading to lower clearance of lipoprotein from our body through LDLR [15]. Other studies have also supported that apoE ϵ 4 allele is associated with the risk of CAD [16]. However, according to a recent meta-analysis, the cardiovascular role of apoE2 is uncertain [17] because of its tendency to increase triglyceride (TG) level [18]. In addition, apo ϵ 2 homozygote in combination with certain additional disorders may develop type III familial hyperlipidemia and premature atherosclerosis [12]. Consequently, with their ability to affect lipid levels, the apoE gene polymorphism could be one of the factors influencing development of both T2DM and CAD. The prevalence of T2DM and CAD is increasing in Thai population according to the cross-sectional ECG survey of 1991 in Thai population, which found that the age-standardized prevalence rate of CAD was 9.9/1000 subjects (men 9.2/1000, women 10.7/1000) [19]. World Health Organization (WHO) global prevalence of diabetes report estimated that by the year 2030, 366 million people, particularly in developing countries, will be affected by diabetes [20]. Genetic factors like apoE are

also considered to be genetic determinants of plasma lipoprotein levels and play a central role in the development of CAD. However, it is unclear whether apoE gene is associated with T2DM. The primary aim of this study was to demonstrate the influence of the apoE gene polymorphism on plasma lipids is notable and is an important determinant of T2DM and CAD. Secondly, this genetic study might also add more information about a Thai population beyond traditional risk factors.

Materials and methods

Subjects

The studied subjects were recruited from Siriraj Hospital in Bangkok province. Full consent forms were signed by the subjects after the nature and motif of the study was clearly explained to them. The study protocol was approved by the Ethics Committee of Clinical Study in Humans, Faculty of Medicine Siriraj Hospital, Mahidol University. Questionnaires were used to collect the information of family and medical history, alcohol consumption, smoking habits and physical activity. Other clinical and biochemical data such as dyslipidemia, systolic blood pressure (SBP) and diastolic blood pressure (DBP) from all subjects were obtained from clinical and laboratory examinations. Anthropometric data (weight, height) were collected and used for BMI calculation. Obesity was defined as BMI ≥ 25 kg/m² according to WHO suggested criteria for Asian populations [21]. Dyslipidemic or hyperlipidemic were defined as when one has level of TC >200 mg/dL, TG >150 mg/dL, LDL-C >130 mg/dL, HDL-C <40 mg/dL, TC/HDL-C ratio >4.0 or under medication of lipid lowering drugs. Cigarette smokers were allowed into the study if they had once smoked even if they were no longer smokers. Alcohol drinkers were defined as those who drank at least two times a week for more than a year. Physical activities were defined as exercise for at least 2 to 3 days/week for at least 30 minutes. Hypertension was defined as blood pressure above 140/90 mmHg or taking antihypertensive drugs. These subjects were categorized into three groups: normal healthy controls, T2DM with and T2DM without CAD, according to the criteria of American Diabetes Association Classification 2010 [22], with an age range from 40-65 years.

The normal healthy control group consisted of 149 subjects with fasting plasma glucose (FPG) < 100 mg/dL from among those who were randomly selected after routine health check-up to screen out those having hyperlipidemia, hypertension, history of chest pain, family history of CAD or any forms of cardiovascular disease, diabetes mellitus, hepatic and renal diseases, inflammation, general illness, traumatic injury, endocrine disease, and other metabolic disorders. Subjects under medication or drug abusers were also excluded.

T2DM subjects without CAD included 155 subjects. All subjects fulfilled the diabetes mellitus diagnostic criteria of FPG ≥ 126 mg/dL or were under treatment with oral antidiabetic drugs. The subjects had no electrocardiogram (ECG) and/or angiography abnormalities, no documented history of CAD and no sign of myocardial ischemia during exercise. Other exclusion criteria for this group were possession of type 1 diabetes mellitus, renal disease, hepatic disease, endocrine disease, and other metabolic diseases.

T2DM complicated with CAD (T2DM + CAD) subjects included 147 subjects. Subjects were confirmed CAD by coronary angiography, with at least 50% stenosis in a major coronary artery or one of their branches. Subjects were diagnosed to have diabetes with FPG ≥ 126 mg/dL or those with drug-treated T2DM. Exclusion criteria included those having renal disease, hepatic disease, type 1 diabetes mellitus, any form of endocrine disease or metabolic disease.

APOE Genotyping

Blood samples were collected in EDTA containing tubes. Guanidine-HCl precipitation method was performed for genomic DNA extraction. Genomic DNA was subjected to polymerase chain reaction (PCR) with primers specific to apoE gene, sense: 5' AACAACTGACCCCGGTGGCG 3', antisense: 5' ATGGCGCTGAGGCCGCGTC 3', sense: 5' CCCACCTGCGCAAGCTGCGC 3', using thermal cycler with thermal profile according to Richard et al [23]. In brief, PCR reaction mixture included 20 pmol of each primer, 0.3 μ g genomic DNA, 10 mM of each dNTP, 10X PCR buffer, 10% DMSO in a final volume of 50 μ L. 10 μ L of PCR products was digested with 0.3 unit of HhaI enzyme according to the supplier's recommended procedure (Biolabs New England). The resulted fragments were then separated on 8% polyacrylamide gel and stained with ethidium bromide. Bands were compared with 25 bp DNA marker and the different individual genotypes were separated and categorized based on the following band length criteria: E2/E2: 91, 83, 61; E3/E3: 91, 61, 48, 35; E4/E4: 72, 61, 48, 35; E2/E3: 91, 83, 61, 48, 35; E2/E4: 91, 83, 72, 61, 48, 35 and E3/E4: 91, 72, 61, 48, 35.

Lipid analysis and biochemical determination

Venous blood samples were collected from the patients after 12 hours of overnight fast. Plasma (TC), TG, HDL-C, glucose and glycosylated hemoglobin (HbA1c) were quantified using an automated clinical chemistry analyzer and enzyme-based colorimetric lists supplied by Roche Diagnostics, Germany. LDL-C levels were calculated using Friedewald formula [24]. VLDL-C level was obtained by using the following equation, VLDL-C = (TC - LDL-C - HDL-C) and non-HDL-C level was

calculated by subtracting HDL-C value from total cholesterol value as a candidate biometrical equivalent to apoB 100 in diabetes [25].

Statistical analysis

All statistical analyses were performed with IBM SPSS v1 (IBM corporation, Armonk, New York, US) and Microsoft 2010 based Excel (Microsoft, US). All data were expressed as mean (SEM). One-way analysis of variance (ANOVA) followed by posthoc Bonferroni multiple comparison test was applied to evaluate the mean difference of the data between three groups (control, T2DM without CAD, and T2DM with CAD). Independent samples t-tests were applied to evaluate the mean difference in HbA1c, diabetes duration between T2DM and T2DM with CAD and lipid profiles between groups. Categorical data such as sex, hypertension, history of smoking, history of alcohol drinking, physical activity and dyslipidemia were evaluated by Chi-square tests or Fisher's exact tests. Allele and genotype difference between groups and deviations from Hardy-Weinberg equilibrium were tested by Chi-square tests. The association between diseases and polymorphism was provided by crude or univariable logistic regression analysis with unadjusted odds ratio (OR) and 95% confidence interval (95% CI). The adjusted OR with 95% CI was used to determine the independent risk factor for development of diabetes and CAD by multivariable logistic regression analysis after adjusting for age, sex, BMI, smoking habits, and physical activity. The multivariable logistic regression analysis was also repeated for adjusted OR with 95% CI to determine the risk of apoE polymorphism combined with obesity and smoking (joint association) after adjusting for age, sex, and physical activity. Statistical significance was considered as $p < 0.05$.

Results

The baseline characteristics of the study population

All enrolled subjects were Thais recruited from Bangkok area. The anthropometric and demographic data were summarized along with the clinical and biochemical data as shown in Table 1. The data from normal controls were used to compare with the data from T2DM with and without CAD. A significant age difference was found between control and T2DM with CAD groups ($p < 0.0001$) which particularly signifies that increase in age could possibly lead to higher chances of developing CAD. There was also significant sex difference between controls and CAD ($p < 0.0001$) due to the high prevalence of males among CAD patients. Other clinical data such as BMI, systolic blood pressure (SBP) and diastolic blood pressure (DBP) were significantly higher in both groups when compared to controls. CAD patients had higher frequencies of smoking ($p = 0.033$) and lower

levels of physical activity ($p = 0.030$) as compared to controls and T2DM patients. Lipid profile data demonstrated significantly higher levels of TC, LDL-C, VLDL-C, TG, non-HDL-C and lower level of HDL-C in T2DM and CAD patients when compared to controls. Patients with CAD had diabetes for a longer period ($p < 0.0001$) and showed no significant difference in HbA1c ($p = 0.151$) as compared to the patients without CAD. Blood glucose levels were also significantly higher in both groups as compared to the controls ($p < 0.0001$).

ApoE genotype and allele frequencies

The genotype distribution of both controls and CAD patients were in Hardy-Weinberg equilibrium except for T2DM that deviated from the basic norm ($p = 0.0001$). This was due to the lower presence of E2/E3 genotype in diabetic group (Table 2). However, since it was unlikely that any of the assumptions for Hardy-Weinberg equilibrium were violated, such a departure was attributed to chance. E3/E3 is also the most common genotype in Thai general population. E3/E3 genotype and $\epsilon 3$ allele were significantly lower ($p = 0.003$ and 0.010 respectively) while E3/E4 genotype was significantly higher only in T2DM with CAD subjects ($p = 0.0004$). In T2DM subjects, E3/E4 genotype had a tendency to be higher (19.35% vs. 14.09%) but showed no significant difference ($p = 0.219$). In addition, E2/E3 genotype and $\epsilon 2$ allele were also significantly lower in T2DM patients ($p = 0.005$ and 0.010 , respectively). However, $\epsilon 4$ allele frequency manifested itself as significantly higher in both T2DM and CAD patients as compared to the controls ($p = 0.047$ and 0.009 , respectively) (Table 3).

Association of apoE gene polymorphism and diseases

Low lipid clearance property of $\epsilon 4$ allele may possibly make it an independent risk factor for the development of CAD and T2DM. Thus, E3/E4 genotype may be the risk factor for development of CAD and/or diabetes. The univariate analysis was used to determine these associations according to the genotype and allele frequencies of apoE gene polymorphism.

Interestingly, E3/E4 genotype increased the risk of CAD with unadjusted OR 2.78 (95%CI 1.50-5.16, $p = 0.0004$) and showed no association with T2DM with unadjusted OR 1.46 (95%CI 0.76-2.81, $p = 0.219$). The $\epsilon 4$ allele appeared to increase risk of both T2DM and CAD with unadjusted OR 1.72 (95%CI 0.97-3.06, $p = 0.047$) and 2.37 (95%CI 1.36-4.15, $p = 0.0009$, respectively) (Table 4). After being adjusted for age, sex, BMI, smoking habits, and physical activity using multivariable binary logistic regression analysis as shown in Table 5 the E3/E4 genotype appeared to be the independent risk factor for development of CAD with adjusted OR 2.52 (95%CI 1.28-4.97, $p = 0.008$). However, the E3/E4 genotype was

Table 1 Anthropometric, demographic and clinical data of T2DM with and without CAD compared to normal healthy controls

Variables	Controls (n = 149)	T2DM (n = 155)	T2DM + CAD (n = 147)
Age (Years)	52.01 (0.62)	51.95 (0.53)	57.56 (0.47)**
Sex (male/female)	49/100	57/98	95/52**
BMI (kg/m ²)	23.84 (0.27)	26.89 (0.31)*	27.49 (0.36)**
SBP (mmHg)	111.54 (1.12)	147.31 (1.76)*	155.46 (1.96)**
DBP (mmHg)	73.11 (0.86)	84.50 (1.01)*	92.26 (1.20)**
Hypertension (%)	-	73.54	93.19
Smokers (%)	3.35	4.51	9.52 [†]
Alcohol consumer (%)	23.80	25.50	32.70
Physical activity (%)	68.70	67.09	56.46 [†]
Diabetes duration (Years)	-	6.09 (0.37)	9.07 (0.62)
Glucose (mg/dL)	89.7 (0.64)	225.9 (6.41)*	192.9 (6.55)**
HbA1c (%)	-	8.94 (0.16)	8.60 (0.17)
Triglyceride (mg/dL)	98.23 (3.68)	221.1 (10.35)*	204.3 (8.70)**
TC (mg/dL)	191.52 (1.80)	237.76 (5.47)*	202.48(4.19) [†]
LDL-C (mg/dL)	111.31(1.76)	146.65(4.84)*	122.36(3.93) [†]
HDL-C (mg/dL)	60.17 (1.23)	50.42 (1.27)*	44.59 (0.95)**
VLDL-C (mg/dL)	19.65 (0.73)	45.86 (2.67)*	40.86 (1.74)**
Non-HDL-C (mg/dL)	131.35 (1.95)	182.26 (4.46)*	157.88 (4.09)**
TC/HDL-C	3.4 (0.1)	4.9 (0.2)*	4.8 (0.1)**
Dyslipidemia (%)	-	85.2	100

Data are presented as mean values (SEM), or numbers (n) and percentage of subjects. *p-value < 0.0001 in comparison between controls and T2DM, **p-value < 0.0001 in comparison between controls and T2DM + CAD, [†]p-value < 0.05 in comparison between controls and T2DM + CAD. BMI: body mass index; SBP: systolic blood pressure; DBP: diastolic blood pressure; HbA1c: hemoglobin A1c; TC: total cholesterol; LDL-C: low density lipoprotein cholesterol; TG: triglyceride; HDL-C: high density lipoprotein cholesterol; VLDL-C: very low density lipoprotein cholesterol.

not found to be an independent risk factor for diabetes, but the ε4 allele was the independent risk factor of both T2DM and CAD with adjusted OR 2.04 (95%CI 1.07-3.86, p = 0.029) and OR 2.32 (95%CI 1.17-4.61, p = 0.016), respectively. The multivariable analysis of apoE4 polymorphism and the risk of T2DM and CAD were also determined according to anthropometric and demographic characteristics. Sex (particularly male gender), age and BMI were also independent risks for CAD but only BMI was the independent risk for T2DM (Table 5).

From joint association analysis, E3/E4 genotype was found to further increase the risk for development of diabetes and CAD when combined with either obesity or smoking after being adjusted for age, sex, and physical activity. The risk factor for T2DM increased from adjusted OR 1.42 (95%CI 0.72-2.78, p = 0.310) (Table 5) to adjusted OR 4.93 (95%CI 1.74-13.98, p = 0.003) (Table 6), whereas CAD risk increased from adjusted OR 2.52 (95%CI 1.28-4.97, p = 0.008) (Table 5) to adjusted OR 10.48 (95%CI 3.56-30.79, P < 0.0001) (Table 6). However, the number of E3/E4

Table 2 Frequency distribution of apoE genotypes and alleles in Hardy-Weinberg Equilibrium

Genotype	Controls (n = 149)	T2DM (n = 155)	T2DM + CAD (n = 147)
E2/E2	2 (1.34%)	1 (0.64%)	1 (0.68%)
E3/E3	113 (75.83%)	117 (75.48%)	88 (59.86%)
E4/E4	1 (0.67%)	4 (2.58%)	1 (0.68%)
E2/E3	12 (8.05%)	2 (1.29%)	11 (7.48%)
E2/E4	0 (0%)	1 (0.64%)	0 (0%)
E3/E4	21 (14.09%)	30 (19.35%)	46 (31.29%)
p-value	0.198	0.0001	0.157
Allele ε2(95% CI)	0.05(0.03 - 0.08)	0.02(0.01 - 0.04)	0.04(0.02 - 0.07)
Allele ε3(95% CI)	0.87(0.82 - 0.90)	0.86(0.81 - 0.89)	0.79 (0.74 - 0.83)
Allele ε4(95% CI)	0.07(0.05 - 0.11)	0.13 (0.09 - 0.16)	0.16 (0.12 - 0.21)

Table 3 Genotype and Allele frequencies distribution of apoE gene polymorphism in controls, T2DM with and without CAD compared to healthy controls

Genotype	Groups		
	Controls (n = 149)	T2DM (n = 155)	T2DM + CAD (n = 147)
E2/E2	0.013	0.006	0.006
E3/E3	0.758	0.754	0.598**
E4/E4	0.007	0.025	0.007
E2/E3	0.080	0.012*	0.074
E2/E4	0	0.006	0
E3/E4	0.140	0.193	0.321***
Allele ε2	0.05	0.02*	0.04
Allele ε3	0.87	0.86	0.79**
Allele ε4	0.07	0.13*	0.16 ^{††}

*p-value < 0.01 compared between controls and T2DM, **p-value < 0.01, ***p-value < 0.001 compared between controls and T2DM + CAD.

genotypes combined with both obesity and smoking in T2DM with and without CAD was small; thus the statistical power of this joint association comparison is limited.

Relationship between Lipid profiles and apoE4-bearing genotypes

The significant differences in apo E distribution among these three groups are mainly due to the differences in frequencies of the ε4 allele. These distributions in apo ε4 allele frequencies may lead to the differences in plasma lipid levels. To test this hypothesis, we analyzed the correlation between apoE4 and plasma lipid levels (Table 7). There were no significant differences between E4 carriers (E2/E4, E3/E4, E4/E4) and E3/E3 genotype as a reference for all values of plasma lipid levels in both control and T2DM with CAD. In T2DM group, there were significant elevation in the values of VLDL-C and TG but there were no significant differences in TC, LDL-C, HDL-C and non-HDL-C levels in E4 carriers. After pooling the overall subjects, the results showed that E4 carrier has significantly elevated VLDL-C and

TG levels while HDL-C concentration was significantly decreased as compared to E3/E3.

Discussion

It has been reported that dyslipidemia or dyslipoproteinemia might strongly contribute towards aggravating the problems of micro- and macroangiopathic complications in diabetic patients [26,27]. This implication was mainly characterized by higher prevalence of male gender, older age, smoking habits, and less physical activity. Other differences were duration of diabetes, lower levels of HDL-C and hypertension when compared to those without CAD which were similar to this study. However, some studies have also shown that elevation of TG and non-esterified fatty acid (NEFA) levels accelerated the pathogenesis of T2DM [28-30]. This suggests that dyslipidemia is associated with both T2DM and CAD. Genetic factors which have strong impact on the metabolism of plasma lipid have been studied regarding the potential effect of T2DM and cardiovascular outcomes in various diabetic and non-diabetic subjects. These studies have underscored that

Table 4 Associations of apoE gene polymorphisms with the risk of T2DM and CAD compared to healthy controls represented as unadjusted OR

Genotype	Controls (n = 149)	T2DM (n = 155)	Unadjusted OR (95% CI)	p-value	T2DM + CAD (n = 147)	Unadjusted OR (95% CI)	p-value
E2/E2	2	1	0.48 (0.02 - 6.78)	0.616	1	0.50 (0.02 - 7.16)	1.000
E3/E3	113	117	0.98 (0.56 - 1.71)	0.942	88	0.48 (0.28 - 0.81)	0.003
E4/E4	1	4	3.92 (0.41 - 93.19)	0.371	1	1.01(0.06 - 16.36)	1.000
E2/E3	12	2	0.15 (0.02 - 0.72)	0.005	11	0.92(0.36 - 2.33)	0.854
E2/E4	0	1	-	1.000	0	-	-
E3/E4	21	30	1.46 (0.76 - 2.81)	0.219	46	2.78(1.50 - 5.16)	0.0004
Allele ε2	16	5	0.29 (0.09 - 0.85)	0.011	13	0.81(0.36 - 1.81)	0.580
Allele ε3	259	266	0.91 (0.56 - 1.48)	0.691	234	0.57(0.36 - 0.90)	0.107
Allele ε4	23	39	1.72 (0.97 - 3.06)	0.047	49	2.37(1.36 - 4.15)	0.0009

Table 5 Association of apoE gene polymorphism with the risk of T2DM and CAD compared to healthy controls represented as adjusted OR

	T2DM		T2DM + CAD	
	Adjusted OR (95% CI) ^{††}	p-value	Adjusted OR (95% CI) ^{††}	p-value
E3/E4 genotype	1.42 (0.72 - 2.78)	0.310	2.52 (1.28 - 4.97)	0.008 ^{††}
ε4 allele	2.04 (1.07 - 3.86)	0.029 ^{††}	2.32 (1.17 - 4.61)	0.016 ^{††}
Age	1.00 (0.97 - 1.04)	0.887	1.13 (1.08 - 1.18)	<0.0001
Sex	1.29 (0.77 - 2.14)	0.326	3.10 (1.75 - 5.49)	<0.0001
BMI	3.68 (2.25 - 6.00)	<0.0001	3.66 (2.09 - 6.39)	<0.0001
Smoking	2.58 (0.87 - 7.66)	0.086	0.59 (0.13 - 2.50)	0.473
Physical activity	0.68 (0.40 - 1.19)	0.148	0.67 (0.36 - 1.24)	0.206

^{††}After adjusting for age, sex, BMI, smoking habit and physical activity. Normal reference ranges of clinical and biochemical data are described in materials and methods.

apoE gene encoding apolipoprotein E is associated with significant variation in lipid profile in our body [31]. ApoE gene is one of the most widely studied candidate genes for CAD or any other form of cardiovascular disease and/or diabetes. A significant relationship of apo E polymorphism and CAD has been observed in several ethnic groups, including Caucasian in the USA [32], Austrian [33], Finnish [34], Italian [35], Turkish [36], Indian [37] and Chinese [38] populations. Some studies have shown apoE ε4 allele as an independent risk factor after further adjustment of other established risk factors for development of CAD in T2DM [16] and myocardial infarction [37] patients. However, no independent association was observed after adjusting for age, sex, smoking, BMI, HDL-C and TG in African- Americans and Caucasians [39]. Therefore, it is of great interest to study the independent risk factors of this gene polymorphism in a Thai population. The present study also coincides with a previous study regarding the development of CAD in T2DM subjects indicating that the frequency of ε4 allele is significantly higher in CAD compared to controls and can be one of the factors for the progression of CAD disease [40]. The frequencies of apoE allele and genotype vary between different populations [13]. In this study, ε3 allele was the most frequent allele in control, diabetes, and CAD subjects, while ε2 was less frequent. E3/E3 genotype was the most frequent isoform found in all groups compared to the other genotypes which corresponds to previous reports

[13,41]. Additionally, the higher frequency of E3/E4 genotype was observed only in CAD (p = 0.0004), whereas the higher frequency of ε4 allele was observed in both T2DM and CAD (p = 0.047, p = 0.0009, respectively). From this study we can conclude that among these selected apoE alleles (ε2, ε3, and ε4), ε4 allele is one of the predictors of both diseases in Thai subjects. In addition, the PDAY (Pathobiological Determinants of Atherosclerosis in Youth) study reported that individuals with E2/E3 genotype had fewer atherosclerotic lesions, whereas those with the E3/E4 had more lesions in the abdominal aorta [42]. These observations strongly suggest that the ε2 allele has a protective role against atherosclerosis. However, in this study, we found that E2/E3 genotype and ε2 allele were also significantly lower in T2DM group indicating that the protective effect of ε2 allele on the development of hyperlipidemia might be less in T2DM patients. It is probable that other environmental and genetic factors are involved in the pathogenesis of CAD.

This present study is the first report to demonstrate that the ε4 allele containing genotypes is the major predictor of development of both T2DM and CAD. Our study also shows strong association of E3/E4 genotype with development of CAD in T2DM patients (Table 4) as reported in previous studies [16,42]. After adjusting for age, sex, smoking, BMI, and physical activity, E3/E4 containing subjects showed 2.52-fold higher risk (p = 0.008) while ε4 allele containing genotypes led to a 2.32-fold

Table 6 Joint association study to evaluate the risk of T2DM and CAD compared to controls

Genotype	Obesity/Smoking	Controls	T2DM	Adjusted OR (95% CI) ^{††}	p-value	T2DM + CAD	Adjusted OR (95% CI) ^{††}	p-value
E3/E3	Neither	70	34	1	-	30	1	-
E3/E3	Either	40	73	3.65 (2.07 - 6.43)	<0.0001	56	2.24 (1.15 - 4.35)	0.018
E3/E3	Both	2	10	10.05 (2.06 - 48.96)	0.004	1	2.02 (0.16 - 25.01)	0.585
E3/E4	Neither	14	14	1.97 (0.84 - 4.64)	0.118	7	1.02 (0.34 - 3.06)	0.970
E3/E4	Either	6	15	4.93 (1.74 - 13.98)	0.003	36	10.48 (3.56 - 30.79)	<0.0001
E3/E4	Both	6	1	-	-	3	-	-

^{††} After adjusting for age, sex and physical activity. Normal reference ranges of clinical and biochemical data are described in materials and methods.

Table 7 Fasting lipid concentration according to apoE4-bearing genotypes

Variables (mg/dL)	Control		T2DM		T2DM + CAD		Total	
	E3/E3 (n = 113)	E4-bearing genotypes (n = 22)	E3/E3 (n = 117)	E4-bearing genotypes (n = 35)	E3/E3 (n = 88)	E4-bearing genotypes (n = 47)	E3/E3 (n = 318)	E4-bearing genotypes (n = 104)
TC	191.58 (2.02)	194.09 (5.22)	232.80 (5.33)	234.77 (8.65)	206.52 (5.26)	192.04 (7.96)	210.88 (2.72)	206.86 (5.11)
LDL-C	110.87 (1.97)	118.52 (4.45)	145.14 (4.54)	140.91 (8.43)	125.33 (4.84)	113.80 (7.33)	127.48 (2.39)	123.92 (4.59)
HDL-C	60.07 (1.43)	55.77 (2.51)	50.84 (1.13)	48.63 (2.98)	45.49 (1.27)	42.70 (1.37)	52.64 (0.81)	47.46 (1.37)**
VLDL-C	20.04 (0.87)	20.18 (1.69)	41.38 (1.99)	55.77 (5.85)*	41.25 (2.37)	38.91 (3.04)	33.76 (1.17)	40.62 (2.72)**
TG	100.20 (4.33)	100.91 (8.48)	206.92 (9.99)	278.83 (29.63)*	206.26 (11.78)	194.53 (15.21)	168.82 (5.88)	203.10 (13.62)**
Non-HDL-C	131.50(2.16)	138.32(4.99)	181.97(5.36)	186.14(8.12)	161.03(5.24)	149.34(7.44)	158.24 (2.83)	159.39 (4.82)

E4-bearing genotypes = E3/E4, E4/E4, E2/E4. Independent sample t-test test was applied to compare between E3/E3 and E4-bearing genotypes. Data are presented as mean values (SEM).

*p-value < 0.05 in comparison between E3/E3 and E4-bearing genotypes in T2DM.

**p-value < 0.01 in comparison between E3/E3 and E4-bearing genotypes in total subjects.

higher risk ($p = 0.016$) for developing CAD and $\epsilon 4$ allele containing genotypes led to a 2.04-fold higher risk ($p = 0.029$) in the development of T2DM as compared to the controls. This signifies that other risk factors have no influence in development or curbing the disease, suggesting that $\epsilon 4$ allele is the independent risk factor for both T2DM and CAD. However, E3/E4 genotype further increases the risk for the development of diabetes and CAD when combined with smoking and/or obesity (Table 6). This confirms the findings that progression and development of diabetes and CAD is the consequence of multifactorial parameters, for example, obesity, smoking, oxidative stress, defect in pancreatic β -cells, alcohol consumption, exercise, hypertension and genetic factors. In addition, incidence of T2DM is also subject to gene-environment interactions, for example, dietary factors, intake of vegetable fat, polyunsaturated fatty acid, dietary fiber (particularly cereal fiber), magnesium, and caffeine were significantly inversely correlated and intakes of trans fat, saturated fatty acid and heme-iron, glycemic index, and glycemic load were significantly positively correlated with the incidence of T2DM [43]. Similarly, regular exercise has been shown to reduce weight, BMI, HDL-C and insulin resistance. This indicates that exercise is also associated with decrease risk of T2DM and CAD. Furthermore, apoE gene polymorphism has been shown to modulate the effects of exercise on lipoprotein concentrations in plasma [44]. In this study, smoking and obesity are shown to be the two major contributing factors that can promote the development of T2DM and CAD or both when placed in joint association (Table 6). The reason for this is that smoking and obesity enhance the oxidative stress which results in decreased insulin secretion from pancreatic β -cell and decreased uptake of blood glucose into the muscle cells [9,10]. This evidence was supported by the experimental study on atherosclerosis-susceptible B6 (B6.apoE $-/-$) and atherosclerosis-resistant BALB (BALB.apoE $-/-$) mice that showed defects in insulin secretion rather than defects in insulin resistance which explains the mark difference in susceptibility to T2DM [45]. Moreover, oxidative stress also increases vascular inflammation leading to CAD and/or any form of cardiovascular disease with or without combination of T2DM [46].

To demonstrate that apoE gene polymorphism is involved in the lipid clearance process and it has great influence on the lipid level in our body [47,48]. Although previous study has shown that non-HDL-C concentration is similar to or better than LDL-C alone in predicting cardiovascular disease (CVD) incidence [49], in this study, the results showed that the decrease in HDL-C and elevation of TG, VLDL-C and LDL-C levels (Table 1) are also the landmarks of diabetes and CAD development as found in an earlier study [50]. However, the association of apoE $\epsilon 4$

allele and lipid profile is still controversial [51]. The $\epsilon 4$ allele has been shown to be associated with high concentration of serum TC and LDL-C in Chinese population [38]. Some studies also showed a significant relationship between E3/E4 genotype with lower HDL-C and higher LDL-C concentrations in CAD patients [52] and with higher TG levels in T2DM patients [48] as compared to the healthy controls. However, a study in the Tunisian population has suggested that HDL-C concentration and other lipid profiles are not associated with apoE gene polymorphism in the total population studied, but has proven that $\epsilon 4$ allele increased LDL-C in type 2 diabetic men [53]. Association of $\epsilon 4$ allele with higher LDL-C and lower HDL-C levels was only found in type 2 diabetic women in Spanish population [54] indicating that gender affects the effect of apoE gene polymorphism. In this study, there is a mean comparison between E4 carriers and E3/E3 genotype in the T2DM and total population studied which showed a significantly higher VLDL-C, TG and lower HDL-C levels in E4 carriers (Table 7) similar to the report of Knouff et al 1999 [15]. Nevertheless, the differences of lipid profiles were not observed in controls and T2DM with CAD. This may be due to dietary restrictions in the controls and lipid-lowering aggressive treatment in T2DM with CAD. Kolovou et al. 2006 [55] reported that the $\epsilon 4$ allele can increase LDL-C concentrations in the presence of an atherogenic diet, but a lower fat diet can suppress this effect. On the other hand, aerobically trained individuals have high HDL and display enhanced glucose tolerance [56]. A lipid-lowering-drug which raises HDL levels and decrease TG levels delays the onset of T2DM and reduces the development of atherosclerosis [57]. In addition, loss of caspase-1 activity which involves the inflammatory process in human atherosclerotic vessel has shown to reduce atherosclerosis lesion formation in Casp1 $^{-/-}$ Apo E $^{-/-}$ mice [58]. Another reason for these variations of lipid profiles might be because of the differences in genetic background and the prevalence of high oxidative stress in the studied population. When considering the basis of gene-gene and gene-environment interactions, as described above, the polymorphism study at the genome level might not provide a real picture for development of diseases in respect to lipid profile. A more robust parallel study at the protein level, or of other candidate genes, may be necessary to evaluate the mechanism of T2DM and CAD development. However, this study will provide a good starting point for the screening of large populations before proceeding to the protein level study of apoE and other candidate genes in various races and, additionally, including pharmacogenomics study in the future.

Conclusions

This study tentatively supports the fact that variability in apoE gene locus is associated with diabetes with and

without CAD complication by influencing the plasma lipid levels that are important risk factors for both T2DM and CAD. No single factor can give a satisfactory explanation regarding development of diabetes and CAD as both diseases are complex diseases. Focusing, therefore, on only one risk factor may be less than optimal in enabling researchers to predict who will develop untoward events in complex diseases like diabetes and CAD. Nonetheless, our study strongly supported that the apoE ε4 allele is an independent risk factor for development of both T2DM and CAD. Moreover, obesity and/or smoking, conditions associated with high oxidative stress, can aggravate the progression of both diseases. Genetic studies can thus provide information that may help to improve the ability to identify individuals, families and populations at increased risk, as well as to improve the clinical management of patients with T2DM and CAD. Such information may be useful in developing public health programs reinforcing primary and secondary prevention for T2DM and CAD. Furthermore, T2DM and CAD patients identified to carry high risk genotype or allele should be treated aggressively to prevent the progression of disease.

Abbreviations

APOE: Apolipoprotein E; CAD: Coronary artery disease; T2DM: Type 2 diabetes mellitus; PCR-RFLP: Polymerase chain reaction-restriction fragment length polymorphism; HDL-C: High-density lipoprotein cholesterol; VLDL-C: Very low density lipoprotein cholesterol; LDL-C: Low-density lipoprotein cholesterol; TC: Total cholesterol; TG: Triglyceride; non-HDL-C: Non-high-density lipoprotein cholesterol; SNPs: Single nucleotide polymorphisms; NF-κB: Nuclear factor kappa B; LDLR: Low density lipoprotein receptor; T: Thymine; C: Cytosine; ECG: Electrocardiogram; WHO: World health organization; SBP: Systolic blood pressure; DBP: Diastolic blood pressure; BMI: Body mass index; FPG: Fasting plasma glucose; Guanidine-HCL: Guanidine-hydrochloride; APO B-100: Apolipoprotein B-100; HbA1c: Glycated hemoglobin A1c; NEFA: Non-esterified fatty acid; PDAY: Pathobiological determinants of atherosclerosis in youth.

Acknowledgements

This study was supported by the Graduate Thesis Scholarship grant from Faculty of Medicine Siriraj Hospital, Mahidol University. The authors wish to thank Dr. Saowalak Hunnangkul for excellent statistical advice and Dr. William M. Honsa for English language editorial support.

Author details

¹Department of Biochemistry, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok 10700, Thailand. ²Department of Medicine, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok 10700, Thailand. ³Department of Physiology, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok 10700, Thailand. ⁴Department of Preventive and Social Medicine, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok 10700, Thailand. ⁵Her Majesty's Cardiac Center, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok 10700, Thailand.

Authors' contributions

All authors fulfill the criteria for authorship. RC carried out the SNPs analysis. AL conceived all of the study and coordination. RC and AL conducted statistical analysis and drafted manuscript. TP, DT, SR, and CS provided the samples for SNP analysis. AL, TP, DT, SS, and CS participated in study conception and design, interpretation of data and critical revision of manuscript for important intellectual content. All authors read and approved the final version of the manuscripts.

Competing interests

The authors declare that they have no competing interests.

Received: 26 March 2012 Accepted: 23 April 2012

Published: 23 April 2012

References

1. King H, Aubert RE, Herman WH: Global burden of diabetes, 1995-2025: prevalence, numerical estimates, and projections. *Diabetes Care* 1998, **21**:1414-1431.
2. Aekplakorn W, Stolk RP, Neal B, Suriyawongpaisal P, Chongsuvivatwong V, Cheepudomwit S, Woodward M: The prevalence and management of diabetes in Thai adults: the international collaborative study of cardiovascular disease in Asia. *Diabetes Care* 2003, **26**:2758-2763.
3. Zhang SH, Reddick RL, Piedrahita JA, Maeda N: Spontaneous hypercholesterolemia and arterial lesions in mice lacking apolipoprotein E. *Science* 1992, **258**:468-471.
4. Plump AS, Smith JD, Hayek T, Aalto-Setälä K, Walsh A, Verstuyft JG, Rubin EM, Breslow JL: Severe hypercholesterolemia and atherosclerosis in apolipoprotein E-deficient mice created by homologous recombination in ES cells. *Cell* 1992, **71**:343-353.
5. Li Y, Li X, Shi L, Yang M, Yang Y, Tao W, Xiong Y, Zhang Y, Yao Y: Association of adiponectin SNP + 45 and SNP + 276 with type 2 diabetes in Han Chinese populations: a meta-analysis of 26 case-control studies. *PLoS One* 2011, **6**:e19686.
6. Zeljko HM, Skaric-Juric T, Narancic NS, Tomas Z, Basic A, Salihovic MP, Starcevic B, Janicijevic B: E2 allele of the apolipoprotein E gene polymorphism is predictive for obesity status in Roma minority population of Croatia. *Lipids Health Dis* 2011, **10**:9.
7. Furukawa S, Fujita T, Shimabukuro M, Iwaki M, Yamada Y, Nakajima Y, Nakayama O, Makishima M, Matsuda M, Shimomura I: Increased oxidative stress in obesity and its impact on metabolic syndrome. *J Clin Invest* 2004, **114**:1752-1761.
8. Boden G, Shulman GI: Free fatty acids in obesity and type 2 diabetes: defining their role in the development of insulin resistance and beta-cell dysfunction. *Eur J Clin Invest* 2002, **32**(Suppl 3):14-23.
9. Martyn JA, Kaneki M, Yasuhara S: Obesity-induced insulin resistance and hyperglycemia: etiologic factors and molecular mechanisms. *Anesthesiology* 2008, **109**:137-148.
10. Evans JL, Goldfine ID, Maddux BA, Grodsky GM: Oxidative stress and stress-activated signaling pathways: a unifying hypothesis of type 2 diabetes. *Endocr Rev* 2002, **23**:599-622.
11. Grundy SM: Drug therapy of the metabolic syndrome: minimizing the emerging crisis in polypharmacy. *Nat Rev Drug Discov* 2006, **5**:295-309.
12. Mahley RW, Rall SC Jr: Apolipoprotein E: far more than a lipid transport protein. *Annu Rev Genomics Hum Genet* 2000, **1**:507-537.
13. Singh PP, Singh M, Mastana SS: APOE distribution in world populations with new data from India and the UK. *Ann Hum Biol* 2006, **33**:279-308.
14. Siest G, Pillot T, Regis-Bailly A, Leininger-Muller B, Steinmetz J, Galteau MM, Visvikis S: Apolipoprotein E: an important gene and protein to follow in laboratory medicine. *Clin Chem* 1995, **41**:1068-1086.
15. Knouff C, Hinsdale ME, Mezdour H, Altenburg MK, Watanabe M, Quarfordt SH, Sullivan PM, Maeda N: Apo E structure determines VLDL clearance and atherosclerosis risk in mice. *J Clin Invest* 1999, **103**:1579-1586.
16. Guang-da X, You-ying L, Zhi-song C, Yu-sheng H, Xiang-jiu Y: Apolipoprotein e4 allele is predictor of coronary artery disease death in elderly patients with type 2 diabetes mellitus. *Atherosclerosis* 2004, **175**:77-81.
17. Wilson PW, Schaefer EJ, Larson MG, Ordovas JM: Apolipoprotein E alleles and risk of coronary disease. A meta-analysis. *Arterioscler Thromb Vasc Biol* 1996, **16**:1250-1255.
18. Dallongeville J, Lussier-Cacan S, Davignon J: Modulation of plasma triglyceride levels by apoE phenotype: a meta-analysis. *J Lipid Res* 1992, **33**:447-454.
19. Tatsananivat P, Klungboonkrong V, Chirawatkul A, Bhuripanyo K, Manmontri A, Chitanondh H, Yipintsoi T: Prevalence of coronary heart disease and major cardiovascular risk factors in Thailand. *Int J Epidemiol* 1998, **27**:405-409.

20. Wild S, Roglic G, Green A, Sicree R, King H: **Global prevalence of diabetes: estimates for the year 2000 and projections for 2030.** *Diabetes Care* 2004, **27**:1047-1053.
21. **The Asia-Pacific Perspective: Redefining obesity and its treatment.** World Health Organization: WHO western pacific region; 2000, 1-56.
22. American Diabetes Association: **Diagnosis and classification of diabetes mellitus.** *Diabetes Care* 2010, **33**(Suppl 1):S62-S69.
23. Richard P, Thomas G, de Zulueta MP, De Gennes JL, Thomas M, Cassaigne A, Bereziat G, Iron A: **Common and rare genotypes of human apolipoprotein E determined by specific restriction profiles of polymerase chain reaction-amplified DNA.** *Clin Chem* 1994, **40**:24-29.
24. Friedewald WT, Levy RI, Fredrickson DS: **Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge.** *Clin Chem* 1972, **18**:499-502.
25. Hermans MP, Sacks FM, Ahn SA, Rousseau MF: **Non-HDL-cholesterol as valid surrogate for apolipoprotein B100 measurement in diabetes: Discriminant Ratio and unbiased equivalence.** *Cardiovasc Diabetol* 2011, **10**:20.
26. Krauss RM, Siri PW: **Dyslipidemia in type 2 diabetes.** *Med Clin North Am* 2004, **88**:897-909, x.
27. Jenkins AJ, Rowley KG, Lyons TJ, Best JD, Hill MA, Klein RL: **Lipoproteins and diabetic microvascular complications.** *Curr Pharm Des* 2004, **10**:3395-3418.
28. Bitzur R, Cohen H, Kamari Y, Shaish A, Harats D: **Triglycerides and HDL cholesterol: stars or second leads in diabetes?** *Diabetes Care* 2009, **32**(Suppl 2):S373-S377.
29. DeFronzo RA: **Insulin resistance, lipotoxicity, type 2 diabetes and atherosclerosis: the missing links. The Claude Bernard Lecture 2009.** *Diabetologia* 2010, **53**:1270-1287.
30. Mooradian AD: **Dyslipidemia in type 2 diabetes mellitus.** *Nat Clin Pract Endocrinol Metab* 2009, **5**:150-159.
31. Eichner JE, Dunn ST, Perveen G, Thompson DM, Stewart KE, Stroehla BC: **Apolipoprotein E polymorphism and cardiovascular disease: a HUGe review.** *Am J Epidemiol* 2002, **155**:487-495.
32. Eichner JE, Kuller LH, Orchard TJ, Grandits GA, McCallum LM, Ferrell RE, Neaton JD: **Relation of apolipoprotein E phenotype to myocardial infarction and mortality from coronary artery disease.** *Am J Cardiol* 1993, **71**:160-165.
33. van Bockxmeer FM, Mamotte CD: **Apolipoprotein epsilon 4 homozygosity in young men with coronary heart disease.** *Lancet* 1992, **340**:879-880.
34. Kuusi T, Nieminen MS, Ehnholm C, Yki-Jarvinen H, Valle M, Nikkila EA, Taskinen MR: **Apoprotein E polymorphism and coronary artery disease. Increased prevalence of apolipoprotein E-4 in angiographically verified coronary patients.** *Arteriosclerosis* 1989, **9**:237-241.
35. Corbo RM, Vilardo T, Ruggeri M, Gemma AT, Scacchi R: **Apolipoprotein E genotype and plasma levels in coronary artery disease. A case-control study in the Italian population.** *Clin Biochem* 1999, **32**:217-222.
36. Attila G, Acarturk E, Eskandari G, Akpinar O, Tuli A, Kanadas IM, Kayrin L: **Effects of apolipoprotein E genotypes and other risk factors on the development of coronary artery disease in Southern Turkey.** *Clinica chimica acta; international journal of clinical chemistry* 2001, **312**:191-196.
37. Kumar P, Luthra K, Dwivedi M, Behl VK, Pandey RM, Misra A: **Apolipoprotein E gene polymorphisms in patients with premature myocardial infarction: a case-controlled study in Asian Indians in North India.** *Ann Clin Biochem* 2003, **40**:382-387.
38. Yan S, Zhou X, Lin Q, Song Y: **Association of polymorphism of apolipoprotein E gene with coronary heart disease in Han Chinese.** *Chin Med J (Engl)* 1999, **112**:224-227.
39. Anuurad E, Yamasaki M, Shachter N, Pearson TA, Berglund L: **ApoE and ApoC-I polymorphisms: association of genotype with cardiovascular disease phenotype in African Americans.** *J Lipid Res* 2009, **50**:1472-1478.
40. Stengard JH, Pekkanen J, Ehnholm C, Nissinen A, Sing CF: **Genotypes with the apolipoprotein epsilon4 allele are predictors of coronary heart disease mortality in a longitudinal study of elderly Finnish men.** *Hum Genet* 1996, **97**:677-684.
41. Kataoka S, Robbins DC, Cowan LD, Go O, Yeh JL, Devereux RB, Fabsitz RR, Lee ET, Welty TK, Howard BV: **Apolipoprotein E polymorphism in American Indians and its relation to plasma lipoproteins and diabetes. The Strong Heart Study.** *Arterioscler Thromb Vasc Biol* 1996, **16**:918-925.
42. McGill HC Jr, McMahan CA: **Determinants of atherosclerosis in the young. Pathobiological Determinants of Atherosclerosis in Youth (PDAY) Research Group.** *Am J Cardiol* 1998, **82**:307-367.
43. Murakami K, Okubo H, Sasaki S: **Effect of dietary factors on incidence of type 2 diabetes: a systematic review of cohort studies.** *J Nutr Sci Vitaminol* 2005, **51**:292-310.
44. Bernstein MS, Costanza MC, James RW, Morris MA, Cambien F, Raoux S, Morabia A: **Physical activity may modulate effects of ApoE genotype on lipid profile.** *Arterioscler Thromb Vasc Biol* 2002, **22**:133-140.
45. Li J, Wang Q, Chai W, Chen MH, Liu Z, Shi W: **Hyperglycemia in apolipoprotein E-deficient mouse strains with different atherosclerosis susceptibility.** *Cardiovasc Diabetol* 2011, **10**:117.
46. Tosukh Wong P, Sangwatanaroj S, Jatuporn S, Prapunwattana P, Saengsiri A, Rattanaprucks S, Srimahachota S, Udayachalerm W, Tangkijvanich P: **The correlation between markers of oxidative stress and risk factors of coronary artery disease in Thai patients.** *Clin Hemorheol Microcirc* 2003, **29**:321-329.
47. Tan CE, Tai ES, Tan CS, Chia KS, Lee J, Chew SK, Ordovas JM: **APOE polymorphism and lipid profile in three ethnic groups in the Singapore population.** *Atherosclerosis* 2003, **170**:253-260.
48. Inamdar PA, Kelkar SM, Devasagayam TP, Bapat MM: **Apolipoprotein E polymorphism in non-insulin-dependent diabetics of Mumbai, India and its effect on plasma lipids and lipoproteins.** *Diabetes Res Clin Pract* 2000, **47**:217-223.
49. Fernandez ML, Webb D: **The LDL to HDL cholesterol ratio as a valuable tool to evaluate coronary heart disease risk.** *J Am Coll Nutr* 2008, **27**:1-5.
50. Washio M, Sasazuki S, Kodama H, Yoshimasu PK, Liu Y, Tanaka K, Tokunaga S, Kono PS, Arai H, Koyanagi S, Hiyamuta K, Doi Y, Kawano MT, Nakagaki MO, Takada K, Nii MT, Shirai K, Ideishi MM, Arakawa MK, Mohri MM, Takeshita A: **Role of hypertension, dyslipidemia and diabetes mellitus in the development of coronary atherosclerosis in Japan.** *Jpn Circ J* 2001, **65**:731-737.
51. Frikke-Schmidt R: **Context-dependent and invariant associations between APOE genotype and levels of lipoproteins and risk of ischemic heart disease: a review.** *Scand J Clin Lab Invest Suppl* 2000, **233**:3-25.
52. Singh PP, Singh M, Bhatnagar DP, Kaur TP, Gaur SK: **Apolipoprotein E polymorphism and its relation to plasma lipids in coronary heart disease.** *Indian J Med Sci* 2008, **62**:105-112.
53. Chaaba R, Attia N, Hammami S, Smaoui M, Ben Hamda K, Mahjoub S, Hammami M: **Association between apolipoprotein E polymorphism, lipids, and coronary artery disease in Tunisian type 2 diabetes.** *J Clin Lipidol* 2008, **2**:360-364.
54. Gomez-Coronado D, Alvarez JJ, Entrala A, Olmos JM, Herrera E, Lasuncion MA: **Apolipoprotein E polymorphism in men and women from a Spanish population: allele frequencies and influence on plasma lipids and apolipoproteins.** *Atherosclerosis* 1999, **147**:167-176.
55. Kolovou GD, Anagnostopoulou KK: **Apolipoprotein E polymorphism, age and coronary heart disease.** *Ageing Res Rev* 2007, **6**:94-108.
56. Kraus WE, Houmard JA, Duscha BD, Knetzger KJ, Wharton MB, McCartney JS, Bales CW, Henes S, Samsa GP, Otvos JD, Kulkarni KR, Slentz CA: **Effects of the amount and intensity of exercise on plasma lipoproteins.** *N Engl J Med* 2002, **347**:1483-1492.
57. Tenenbaum A, Motro M, Fisman EZ, Schwammenthal E, Adler Y, Goldenberg I, Leor J, Boyko V, Mandelzweig L, Behar S: **Peroxisome proliferator-activated receptor ligand bezafibrate for prevention of type 2 diabetes mellitus in patients with coronary artery disease.** *Circulation* 2004, **109**:2197-2202.
58. Gage J, Hasu M, Thabet M, Whitman SC: **Caspase-1 Deficiency Decreases Atherosclerosis in Apolipoprotein E-Null Mice.** *Can J Cardiol* 2012, **28**:222-229.

doi:10.1186/1475-2840-11-36

Cite this article as: Chaudhary et al.: Apolipoprotein E gene polymorphism: effects on plasma lipids and risk of type 2 diabetes and coronary artery disease. *Cardiovascular Diabetology* 2012 **11**:36.