

# PUBLISHED VERSION


Qili Fei, Li Yang, Wanqi Liang, Dabing Zhang, Blake C. Meyers  
**Dynamic changes of small RNAs in rice spikelet development reveal specialized reproductive phasiRNA pathways**  
Journal of Experimental Botany, 2016; 67(21):6037-6049

© The Author 2016. Published by Oxford University Press on behalf of the Society for Experimental Biology. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Originally published at:  
<http://doi.org/10.1093/jxb/erw361>

## PERMISSIONS

<http://creativecommons.org/licenses/by/3.0/>



The image shows a Creative Commons Attribution 3.0 Unported (CC BY 3.0) license banner. It features the CC logo and a person icon in a blue circle. The text reads: "Attribution 3.0 Unported (CC BY 3.0)". Below this, it states: "This is a human-readable summary of (and not a substitute for) the [license](#). [Disclaimer](#)." The banner is divided into two main sections: "You are free to:" and "Under the following terms:". Under "You are free to:", it lists "Share" (copy and redistribute) and "Adapt" (remix, transform, and build upon). A circular seal on the right says "Free Cultural APPROVED FOR Works". Under "Under the following terms:", it lists "Attribution" (give credit, link to license, indicate changes) and "No additional restrictions" (no legal terms or technological measures).

**Attribution 3.0 Unported (CC BY 3.0)**

This is a human-readable summary of (and not a substitute for) the [license](#). [Disclaimer](#).

**You are free to:**

- Share** — copy and redistribute the material in any medium or format
- Adapt** — remix, transform, and build upon the material for any purpose, even commercially.

The licensor cannot revoke these freedoms as long as you follow the [license terms](#).

**Under the following terms:**

- Attribution** — You must give [appropriate credit](#), provide a link to the license, and [indicate if changes were made](#). You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.
- No additional restrictions** — You may not apply legal terms or [technological measures](#) that legally restrict others from doing anything the license permits.

1 November 2017

<http://hdl.handle.net/2440/107095>



RESEARCH PAPER

# Dynamic changes of small RNAs in rice spikelet development reveal specialized reproductive phasiRNA pathways

Qili Fei<sup>1,†</sup>, Li Yang<sup>2,†</sup>, Wanqi Liang<sup>2</sup>, Dabing Zhang<sup>2,3,\*</sup> and Blake C. Meyers<sup>1,4,5,\*</sup>

<sup>1</sup> Department of Plant & Soil Sciences and Delaware Biotechnology Institute, University of Delaware, Newark, DE 19711, USA

<sup>2</sup> State Key Laboratory of Hybrid Rice, Joint International Research Laboratory of Metabolic and Developmental Sciences, Shanghai Jiao Tong University and University of Adelaide Joint Centre for Agriculture and Health, School of Life Sciences and Biotechnology, Shanghai Jiao Tong University, Shanghai 200240, China

<sup>3</sup> School of Agriculture, Food and Wine, University of Adelaide, South Australia 5064, Australia

<sup>4</sup> Donald Danforth Plant Science Center, 975 North Warson Road, St. Louis, MO 63132, USA

<sup>5</sup> University of Missouri – Columbia, Division of Plant Sciences, 52 Agriculture Lab, Columbia, MO 65211, USA

\* Correspondence: [bmeyers@danforthcenter.org](mailto:bmeyers@danforthcenter.org) and [zhangdb@sjtu.edu.cn](mailto:zhangdb@sjtu.edu.cn)

† These authors contributed equally to this work.

Received 1 June 2016; Accepted 5 September 2016

Editor: Zoe Wilson, University of Nottingham

## Abstract

Dissection of the genetic pathways and mechanisms by which anther development occurs in grasses is crucial for both a basic understanding of plant development and for examining traits of agronomic importance such as male sterility. In rice, MULTIPLE SPOROCTES1 (MSP1), a leucine-rich-repeat receptor kinase, plays an important role in anther development by limiting the number of sporocytes. *OsTDL1a* (a *TPD1*-like gene in rice) encodes a small protein that acts as a cofactor of MSP1 in the same regulatory pathway. In this study, we analyzed small RNA and mRNA changes in different stages of spikelets from wild-type rice, and from *msh1* and *ostd1a* mutants. Analysis of the small RNA data identified miRNAs demonstrating differential abundances. miR2275 was depleted in the two rice mutants; this miRNA is specifically enriched in anthers and functions to trigger the production of 24-nt phased secondary siRNAs (phasiRNAs) from *PHAS* loci. We observed that the 24-nt phasiRNAs as well as their precursor *PHAS* mRNAs were also depleted in the two mutants. An analysis of co-expression identified three Argonaute-encoding genes (*OsAGO1d*, *OsAGO2b*, and *OsAGO18*) that accumulate transcripts coordinately with phasiRNAs, suggesting a functional relationship. By mRNA *in situ* analysis, we demonstrated a strong correlation between the spatiotemporal pattern of these *OsAGO* transcripts and phasiRNA accumulations.

**Key words:** Anther, Argonaute, microRNA, phasiRNA, rice, spikelet.

## Introduction

Rice (*Oryza sativa*), as a major crop, has been widely used as a monocot model species to explore the genetic basis of flower development in higher plants (Yoshida and Nagato, 2011). Rice anther development is one of the major topics studied

in rice flower development, and changes in the cytological morphology in different developmental stages of rice anthers have been well-described (Zhang and Wilson, 2009; Zhang *et al.*, 2011). Tapetum and microsporocyte specification is

a crucial event in male fertility, occurring at early stages of anther development in plants; a number of genes have been discovered as regulators of cell fate specification (Zhang and Yang, 2014). For example, EXCESS MICROSPOROXYTES 1 (EMS1, or EXTRA SPOROGENOUS CELLS, EXS), a member of the leucine-rich repeat receptor-like kinase (LRR-RLK) family, specifies tapetal identity and limits the number of pollen mother cells (PMCs) in Arabidopsis (Canales *et al.*, 2002; Zhao *et al.*, 2002). The small secreted protein, TAPETAL DETERMINANT 1 (TPD1), has been reported as a ligand of EMS1/EXS with a deterministic role in the cell fate of the tapetum (Jia *et al.*, 2008).

The EMS1/EXS ortholog in rice is MULTIPLE SPOROXYTE (MSP1) (Nonomura *et al.*, 2003), while the ligand protein TPD1 has two TPD1-like orthologs in rice, including OsTDL1A and OsTDL1B, among which OsTDL1A (also known as MICROSPORELESS2, MIL2) may interact with MSP1 (Zhao *et al.*, 2008; Hong *et al.*, 2012). Although *msh1* and *ostd11a* mutants display defects in anther and ovule development, both show a phenotype of complete male sterility, while partially maintaining female fertility (Nonomura *et al.*, 2003; Hong *et al.*, 2012; Yang *et al.*, 2016). MSP1, as a receptor-like kinase in an upstream signaling pathway, affects many other downstream genes involved in rice anther development. For example, a loss of function of MSP1 will largely down-regulate the expression of other genes involved in rice anther development, such as *Undeveloped Tapetum1* (*UDT1*) and *Tapetum Degeneration Retardation* (*TDR*) (Jung *et al.*, 2005; Li *et al.*, 2006). Maize *MAC1* is the ortholog of *OsTDL1A*, having similar functions in limiting archesporial cell proliferation in maize anthers (Wang *et al.*, 2012). Therefore, the OsTDL1A-MSP1 pathway plays a central role in early stages of rice anther development to simultaneously specify the tapetum and limit the number of pollen mother cells (Zhang and Yang, 2014).

Small RNA pathways play roles in both flower development and gametogenesis in plants. Some conserved miRNAs appear to function similarly in flower development across different plant species, such as Arabidopsis, tomato, petunia, rice, and maize (Luo *et al.*, 2013). In rice, miR172 targets *APETALA2* (*AP2*) genes controlling inflorescence architecture and spikelet meristem identity (Zhu *et al.*, 2009; Lee and An, 2012). A number of rice *SQUAMOSA Promoter Binding Protein-Like* (*OsSPL*) genes, including *OsSPL14*, are targeted by miR156; this pathway has a role in flowering time, panicle architecture, grain yield, and other developmental phenotypes (Xie *et al.*, 2006; Jiao *et al.*, 2010; Miura *et al.*, 2010). Other miRNAs, such as miR159 and miR164, are also reported to be involved in rice floral development (Tsuji *et al.*, 2006; Adam *et al.*, 2011). In addition to miRNAs, *trans*-acting siRNAs (tasiRNAs), dependent on the activities of RNA-DEPENDENT RNA POLYMERASE 6 (RDR6) and DICER-LIKE 4 (DCL4), generated from non-coding transcripts such as *TAS3*, play roles in both vegetative and reproductive development in both rice and maize by targeting genes encoding auxin response factors (ARFs). For example, the rice mutant of *SHOOTLESS2* (*SHL2*), the ortholog of Arabidopsis *RDR6*, displays a severe phenotype

of misregulation of adaxial-abaxial polarity patterning in both the lemma and anther (Toriba *et al.*, 2010), while the maize mutant *leafbladeless1* (*lbl1*, a loss of function of an *SGS3* ortholog, *SUPPRESSOR OF GENE SILENCING 3*) is defective in tasiRNA biogenesis and shows a pleiotropic phenotype, including sterile male inflorescences (Nogueira *et al.*, 2007). The RDR6-dependent small RNA biogenesis pathway not only produces tasiRNAs, but also yields two large populations of phased secondary siRNAs (phasiRNAs) in the reproductive tissues of monocots (reviewed in Fei *et al.*, 2013). Data have suggested that these 21- and 24-nt reproductive phasiRNAs, triggered by miR2118 and miR2275, respectively, may play crucial roles in microgametogenesis in maize, because the accumulation of phasiRNAs shows highly stage-specific patterns in maize anther development (Zhai *et al.*, 2015). In addition, the Argonaute (AGO) protein MEIOSIS ARRESTED AT LEPTOTENE1 (*MEL1*), previously demonstrated to be essential for sporogenesis in rice anthers (Nonomura *et al.*, 2007), has recently been shown to recruit 21-nt phasiRNAs (Komiya *et al.*, 2014).

In a recent study, we showed that the MSP1-OsTDL1A partners are master regulators of downstream transcription factors that are involved in plant anther development (Yang *et al.*, 2016). Therefore, we hypothesized that loss-of-function of MSP1-OsTDL1A may also cause great downstream changes in non-coding RNAs and small RNAs. Here, we systematically characterized changes in small RNA and mRNA, especially non-coding *PHAS* transcripts, across early developmental stages of rice spikelets in wild-type, *msh1* and *ostd11a* backgrounds using deep sequencing data. We found comprehensive changes of miRNAs, phasiRNAs, and *PHAS* transcripts in early stages of rice spikelet development. Importantly, the reproductive phasiRNAs displayed stage-specific expression patterns during early stages of anther development, suggesting that the timing of phasiRNA biogenesis is crucial in rice microsporogenesis. Furthermore, phasiRNA and mRNA changes in different developmental stages and mutant backgrounds facilitated the identification of several rice AGOs, in addition to *MEL1*, that potentially load phasiRNAs.

## Materials and methods

### Plant materials and growth conditions

All the rice plants used in this study were in a genetic background of variety 9522, a *japonica* rice. The two male-sterile mutants, *ostd11a* and *msh1-4*, are from a rice mutant library made by <sup>60</sup>Co  $\gamma$ -ray radiation; the molecular details of these mutants are described in Yang *et al.* (2016). Plants were grown in the paddy field of Shanghai Jiao Tong University in China.

### Small RNA and RNA-seq library construction

For small RNA library construction, total RNA enriched for small RNA was extracted. The small RNA fraction between 18 to 30 nt in length was collected by gel separation, then ligated to 5' and 3' adaptors and purified. These small RNAs were reverse transcribed by RT-PCR and finally amplified via PCR. For RNA-seq libraries, after the total RNA extraction and DNase I treatment, magnetic beads with oligo(dT) were used to isolate mRNA. Mixed with the fragmentation

buffer, the mRNA was fragmented into short fragments. The cDNA was synthesized using the mRNA fragments as templates. Short fragments were purified for end repair and single nucleotide A (adenine) addition, ligated to adapters, and then the second strand was degraded using UNG (Uracil-N-Glycosylase). After agarose gel electrophoresis, the suitable fragments were selected for PCR amplification as templates. All the small RNA and RNA-seq libraries were sequenced on an Illumina HiSeq 2000 platform by BGI (BGI-Shenzhen, China).

#### Small RNA data analysis

Small RNA sequencing data were preprocessed by removing adapters, and then mapped to the version 7.0 of the rice genome assembly from the Rice Genome Annotation Project Database (<http://rice.plantbiology.msu.edu>, accessed 23 September 2016) using the program Bowtie (Langmead *et al.*, 2009). Small RNA reads that were mapped to the tRNAs and rRNAs were filtered, and reads mapped to the rice genome from different libraries were normalized to reads per 5 million reads (RP5M) for comparisons. The Bioconductor ([www.bioconductor.org](http://www.bioconductor.org)) package ‘edgeR’ was used for small RNA differential analysis ( $P < 0.05$ , FDR  $< 0.05$ ). Rice miRNA sequences were downloaded from miRBase (Release 21; <http://www.mirbase.org/>, accessed 23 September 2016) (Kozomara and Griffiths-Jones, 2014). The R ([www.r-project.org](http://www.r-project.org), accessed 23 September 2016) package ‘pheatmap’ was used to represent the average abundance of miRNAs from three biological replicates. *PHAS* locus identification was performed using the same method described previously by Zhai *et al.* (2011). Briefly, small RNA sequencing data from different libraries were combined together to increase the sequencing depth for *PHAS* loci identification. A phasing score of 25 was used as a stringent cut-off, followed by a manual check to remove loci producing highly abundant small RNAs in other sizes, which are most likely degradation products from t/rRNAs. The overall phasiRNA abundance for each *PHAS* locus was calculated by summing up the normalized abundance of 21- or 24-nt small RNAs generated from each corresponding 21-*PHAS* and 24-*PHAS* locus.

#### RNA-seq data analysis

Paired-end strand-specific RNA-seq reads (90 bp  $\times$  2) were mapped to the rice genome sequences allowing no more than two mismatches using ‘Tophat’ (Trapnell *et al.*, 2009). The BAM files generated by ‘Tophat’ were sorted and indexed using ‘SAMtools’ (Li *et al.*, 2009), and then visualized via Integrative Genomics Viewer (IGV) (Robinson *et al.*, 2011). The program ‘Cufflinks’ (Trapnell *et al.*, 2012) was used for transcriptome assembly, differential analysis of gene expression, and calculation of the FPKM value (fragments per kilobase of transcript per million mapped reads). For differential analysis of gene expression, we used ‘q-value  $< 0.01$ ’ and ‘fold change  $> 2$ ’ as cut-offs. Bar graphs and line charts representing FPKM values of gene expression were plotted using the Bioconductor package ‘cummeRbund’ (Trapnell *et al.*, 2012).

#### Microarray data analysis

Gene lists were inputted into the webserver Rice Oligonucleotide Array Database (<http://www.ricearray.org/>, accessed 23 September 2016; Cao *et al.*, 2012). Specific public microarray datasets were selected to acquire the abundance values of each gene. The normalized gene expression values were further visualized as heatmaps using the R package ‘pheatmap’.

#### In situ hybridizations

Freshly collected samples were fixed in formalin-acetic acid-alcohol (FAA) and dehydrated in a series of graded ethanol concentrations; these samples were then infiltrated with Histo-clear II, embedded in Paraplast Plus, and subsequently processed into 6- $\mu$ m thick sections using a Leica RM2245 rotary microtome. Templates for RNA probe

synthesis were amplified by PCR from the cDNA. Probes were transcribed *in vitro* under the T7 promoter with RNA polymerase, using the DIG RNA labeling kit (Roche). The RNA *in situ* hybridizations were carried out as described by Kouchi and Hata (1993) and Li *et al.* (2006). The forward and reverse RT-PCR primers were as follows: *OsAGO1d*, 5'-GCAATACCACCCACAAGGAC-3' and 5'-GGTTCCAATACTCCCACTTCC-3'; *OsAGO18*, 5'-CAGTAT AACAGTACGGAACGC-3' and 5'-TGTCATTACAACAAGTAG GAGG-3'.

#### Accession numbers

RNA-seq and small RNA data are available from Genbank, under GEO accession number GSE77300.


## Results

### Comparative analysis of small RNAs in spikelets of wild-type and mutant rice

Small RNAs play crucial roles in mediating both transcription and translation, with different classes distinguishable by their distinct biogenesis pathways (Axtell, 2013). The recent discovery of reproductive phasiRNAs in monocots indicates that this special class of small RNAs may be important for male reproduction, although the underlying mechanism remains to be elucidated (Johnson *et al.*, 2009; Song *et al.*, 2012; Zhai *et al.*, 2015). To assess small RNA and mRNA changes across different stages of rice spikelet development and to understand how they are impacted by perturbation of the OsTDL1A-MSP1 pathway, we prepared small RNA and RNA-seq libraries from spikelets of wild-type rice cultivar 9522, and the mutants *msp1-4* (*msp1* hereafter) and *ostdl1a* [library information is listed in Supplementary Table S1 at JXB online; mutant information is described in Yang *et al.* (2016), and is also shown in Supplementary Fig. S1]. We performed three biological replicates for each genotype and stage. The lengths of rice spikelets correspond to different anther developmental stages (Fig. 1). Specifically, stage 3 (0.15–0.2 mm), stage 5 (0.25–0.3 mm), and stage 7 (0.4–0.45 mm) of rice anthers correspond to 0.5–0.6 mm, 1.0–1.5 mm, and 2.5–3.0 mm rice spikelets, respectively (Zhang *et al.*, 2011) (Fig. 1A); hereafter, we will refer to these sizes of rice spikelets as stage 3, stage 5, and stage 7 spikelets, respectively. Samples were collected at these three stages because MSP1 and OsTDL1A mainly function at early stages (stage 3 to stage 5) of rice anther development (Yang *et al.*, 2016), while stage 5–7 is an important stage at which meiocytes start meiosis (Zhang *et al.*, 2011).

Sequencing reads of small RNAs were aligned to the rice genome and normalized to 5 million (5M), and the distribution of lengths in different stages was analyzed. The three replicates were nearly identical (Fig. 1). The 24-nt small RNAs account for ~75% of all small RNAs at stage 3 in wild-type rice. Interestingly, with the development of spikelets, there was a shift in the predominant size class, with 21-nt small RNAs the largest proportion (nearly 50%) in the profile at stage 5. In stage 7, the proportion (~55%) of 24-nt small RNAs was once again larger than the 21-nt counterpart (<40%), but was far smaller than that in stage 3. The fluctuation of small RNA percentages in different sizes that accompanied





**Fig. 1.** Small RNA size distribution in different developmental stages and backgrounds of rice spikelets. (A) Schematic representation of rice anther structures in different stages of spikelets of wild-type rice. Each layer of cells is indicated by an arrow: epidermis (E); primary parietal cell (PPC); archesporial cell (Ar); endothecium (En); middle layer (ML); tapetum (Ta); sporogenous cell (Sp). Small RNA size distributions in the different stages of wild-type rice spikelets are shown below. (B) Schematic representation of rice anther structures in different stages of spikelets of the *msp1* and *ostd1a* mutants. Excessive archesporial cell (eAr); unknown identity cell (UIC); excessive sporogenous cell (eSp). Small RNA size distributions in the different stages of the mutant spikelets are shown below.

spikelet development may represent shifts in small RNA biogenesis and transcriptome changes in rice reproductive tissues, consistent with prior reports in maize (Zhai *et al.*, 2015). Similar to the wild-type spikelets, in the two mutants changes in small RNA size proportions occurred more significantly in the transition from stage 3 to stage 5 than from stage 5 to stage 7. The only apparent difference was a reduction in the proportion of 21-nt small RNAs in the mutants in stage 5 spikelets (Fig. 1B). To investigate this difference, and others

less readily apparent, we checked the levels of small RNAs that are often 21 nt in length, miRNAs and phasiRNAs.

#### *miRNA expression patterns in different developmental stages of rice spikelets*


A number of miRNAs have characterized roles in plant development, targeting several families of transcription factors or other development-related genes (Jones-Rhoades *et al.*,

2006; Chen, 2009). miRNAs, such as miR156 and miR172, have been proven to control flower development at a post-transcriptional level in both Arabidopsis and rice (Aukerman and Sakai, 2003; Chen, 2004; Xie *et al.*, 2006; Wu *et al.*, 2009a; Zhu *et al.*, 2009; Jiao *et al.*, 2010; Lee and An, 2012). Therefore, the expression patterns of miRNAs are important to our understanding of rice spikelet development.

Genome-wide differential analysis of miRNA expression was performed in wild-type rice across different stages of spikelet development (see Supplementary Tables S2 and S3). From stage 3 to stage 5, >60 miRNAs increased significantly, while only 18 miRNAs decreased during the same time period; from stage 5 to stage 7, the numbers of miRNAs with significantly different levels (up or down) were both fewer than 20 (Fig. 2A; Table 1). Comparing different stages in the *msp1* and *ostd11a* mutant backgrounds, similar sets (both qualitatively and quantitatively) of differentially accumulating miRNAs were observed, suggesting that the male sterile phenotype of these mutants has a limited impact on miRNA levels, relative to the wild-type. This was further confirmed by the observation that very few miRNAs were identified as significantly different in their levels when comparing the two mutants with the wild-type (Table 1). Interestingly, among the few impacted miRNAs in the two mutants, miR2275, which

triggers 24-nt phasiRNA production, was totally abolished (Supplementary Table S3, and see below).

Among the miRNAs differentially expressed in wild-type rice across the developmental stages, the level of miR164 showed a ~7-fold increase from stage 3 to stage 7, whereas miR172 showed a dramatic decline of ~4-fold from stage 3 to stage 5, then remained relatively steady to stage 7 (see Supplementary Fig. S2A). Prior work using PARE/degradome data confirmed the targets of a number of miRNAs in rice (Li *et al.*, 2010; Zhou *et al.*, 2010). From RNA-seq data, we obtained the expression levels of both miR164 and miR172 target genes. Transcript levels of the most abundant miR164 target (LOC\_Os12g05260) decreased from 180 FPKM (stage 3) to 125 FPKM (stage 5), and the expression level increased slightly in stage 7 compared to stage 5, and a similar trend was observed for the target gene LOC\_Os06g23650 (Supplementary Fig. S2B); these were the only two targets that showed an inverse relationship with miR164 abundance. Three miR172 targets showed an inverse correlation with miR172 levels (Supplementary Fig. S2C). miR156 showed an interesting pattern across the developmental stages of rice spikelets: a 21-nt miR156 isoform increased gradually from stage 3 to stage 7, while the 20-nt miR156 firstly decreased



**Fig. 2.** miRNA expression in different developmental stages of spikelets in wild-type rice cultivar 9522. (A) Differentially expressed miRNAs in different developmental stages of spikelets. Only miRNAs with abundance greater than 50 RP5M are included in the heatmap. (B) 20-nt miR156 and 21-nt isoforms display distinct expression patterns; levels indicated as '20/21' show the sum of abundance of the 20 and 21 nt isoforms. (C) Expression levels of miR156 target genes in different stages of rice spikelet development.

**Table 1.** Numbers of differentially abundant miRNAs, identified by pairwise comparisons of rice spikelets in different stages and backgrounds.

	9522_S3	9522_S5	9522_S7	<i>msp1</i> _S3	<i>msp1</i> _S5	<i>msp1</i> _S7	<i>ostdl1a</i> _S3	<i>ostdl1a</i> _S5	<i>ostdl1a</i> _S7
9522_S3	-	63 ↑ 18 ↓	82 ↑ 28 ↓	0 ↑ 1 ↓	-	-	0 ↑ 2 ↓	-	-
9522_S5	-	-	19 ↑ 11 ↓	-	0 ↑ 9 ↓	-	-	2 ↑ 6 ↓	-
9522_S7	-	-	-	-	-	8 ↑ 4 ↓	-	-	13 ↑ 5 ↓
<i>msp1</i> _S3	-	-	-	-	62 ↑ 20 ↓	74 ↑ 26 ↓	-	-	-
<i>msp1</i> _S5	-	-	-	-	-	27 ↑ 16 ↓	-	-	-
<i>msp1</i> _S7	-	-	-	-	-	-	-	-	-
<i>ostdl1a</i> _S3	-	-	-	-	-	-	-	57 ↑ 13 ↓	72 ↑ 14 ↓
<i>ostdl1a</i> _S5	-	-	-	-	-	-	-	-	28 ↑ 3 ↓
<i>ostdl1a</i> _S7	-	-	-	-	-	-	-	-	-

Numbers of significantly up- and down-regulated miRNAs, as indicated by the arrows ( $P < 0.05$ ,  $FDR < 0.05$ ). miRNAs are from the miRBase release 21. 'S3', 'S5', and 'S7' refer to stages 3, 5, and 7, respectively.


from stage 3 to stage 5, and then increased at stage 7 (Fig. 2B). The underlying mechanism is unknown, but it is possible that these two isoforms of miR156 are generated from different *MIR156* genes, and are differentially regulated with distinct targets or cellular expression patterns, exerting distinct patterns of control on the expression of *OsSPLs*. Therefore, studies of individual *MIR156* genes would be important in the future to demonstrate how the miR156 family members differentially control spikelet or panicle development. Similar to miR164 and miR172, a subset of miR156 targets showed an inverse correlation with miR156 (Fig. 2C). Overall, the abundance levels of miRNA target transcripts in the two mutants were quite similar to those in wild-type, indicating that mutations in the OsTDL1A-MSP1 pathway had a limited impact on rice miRNAs and their targets in spikelets (Table 1; Supplementary Fig. S2B–D). Among the miRNA targets validated from prior work, only a subset had the expected inverse correlation with miRNA levels; thus, other factors may impact target transcript levels, including both transcriptional (e.g. transcription factors and epigenetic regulation) and post-transcriptional (e.g. mRNA turnover or sequestration) factors. Taken together, these data indicate that miRNAs together with their target genes are dynamically modulated during early stages of rice spikelet development, and are largely independent of the OsTDL1A-MSP1 pathway.

#### Timing of reproductive phasiRNA biogenesis

PhasiRNAs, in addition to miRNAs, represent another class of small RNAs of great interest, because two populations are exclusively abundant in the reproductive tissues of monocots (Arikiti et al., 2013). More specifically, miR2118 and miR2275 are triggers of 21-nt and 24-nt reproductive phasiRNAs, respectively, in both rice and maize (Song et al., 2012; Zhai et al., 2015). We checked the abundance levels of both miRNAs in different stages and backgrounds; miR2118 abundance peaked at stage 5 in wild-type spikelet libraries, reaching 320 reads per 5 million reads (RP5M), and then

dropped to 130 RP5M at stage 7 (Fig. 3A). Similar abundances were observed in *msp1* and *ostdl1a* mutants, indicating that miR2118 is not impacted in both mutants, possibly explained by miR2118 accumulation in the epidermis of anthers (Zhai et al., 2015), a cell layer apparently not defective in the *msp1* and *ostdl1a* mutants or in the *mac1* maize mutant. In contrast to miR2118, in wild-type spikelets, miR2275 was not expressed in stage 3, and then increased in stage 5 (~85 RP5M) and stage 7 (~240 RP5M) (Fig. 3B); miR2275, as mentioned above, is essentially absent in both *msp1* and *ostdl1a* mutants, at all stages.

We calculated both 21- and 24-nt phasiRNA abundances to see how phasiRNAs change during rice spikelet development and whether they are affected in the two mutants. Phasing analysis resulted in 1843 21-*PHAS* loci and 50 24-*PHAS* loci (see Supplementary Table S4). By summing phasiRNA abundances from each *PHAS* locus, we obtained the overall phasiRNA abundances for both 21- and 24-nt phasiRNAs. In wild-type rice spikelets, the overall abundance of 21-nt phasiRNAs was ~120 000 RP5M in stage 3, increasing by >10-fold in stage 5, and then decreasing by <30% in stage 7 (Fig. 3C). Compared to the wild-type, 21-nt phasiRNAs had a very similar pattern in the two mutants, although with a slightly lower total abundance in stage 5. The 24-nt counterparts had only a few hundred reads in stage 3 and stage 5, but then increased to more than 180 000 RP5M in stage 7 of the wild-type spikelets (Fig. 3D). Consistent with the observation that miR2275 is absent in *msp1* and *ostdl1a* mutants at all stages, 24-nt phasiRNAs were also diminished in both mutants. Considering the developmental defect in these mutants is largely in the anthers and not other tissues in the spikelets (Yang et al., 2016), we infer that the loss of 24-nt phasiRNAs is due to the defective anther development. Furthermore, as in maize (Zhai et al., 2015), rice 21-nt phasiRNAs initiated at an early stage in anther development, the 24-nt phasiRNAs appeared later, coincident with or just before meiosis, and the absence of miR2275 and 24-nt phasiRNAs in *msp1* and *ostdl1a* mutants is similar to maize *mac1* (the ortholog of *ostdl1a* in maize).



**Fig. 3.** miR2118, miR2275, and phasiRNA abundances in rice spikelets. Levels of miR2118 (A), miR2275 (B), miR2118-triggered 21-nt phasiRNAs (C), and miR2275-triggered 24-nt phasiRNAs (D) in different stages and backgrounds of rice spikelets. (E) The percentage of 21-nt (top, blue slices with percentage numbers) and 24-nt phasiRNAs (bottom, orange slices with percentage numbers) out of the total population of 21-nt and 24-nt genome-matched small RNAs in wild-type rice spikelets.

We next calculated the proportion of 21- and 24-nt phasiRNAs in the entire genome-matched populations of the 21- and 24-nt small RNAs in rice spikelets (Fig. 3E). We found that 21-nt phasiRNAs accounted for 24% of total 21-nt siRNAs in wild-type rice spikelets, increased to a remarkable 74% of the total in stage 5, and then reduced slightly to 68% at stage 7. This compares to 60% of all 21-mers at the peak in maize (Zhai *et al.*, 2015); this higher proportion in rice spikelets versus isolated maize anthers may reflect the

fact that there are ~4-fold as many genomic loci generating 21-nt phasiRNAs in rice compared to maize. The 24-nt phasiRNAs were almost absent at stages 3 and 5, followed by a substantial increase to 7% of the total at stage 7 (Fig. 3E). This compares to 64% of the total 24-mers in isolated maize anthers (Zhai *et al.*, 2015); this much lower proportion in rice may reflect the fact that there are fewer loci, or that the peak abundance of 24-nt phasiRNAs is later than stage 7. Overall, these results reveal that both miRNA triggers and



phasiRNAs are largely up-regulated at specific stages in rice anthers.

#### *Interconnected PHAS locus transcription and phasiRNA bursts*

We next examined the phasiRNA precursors (*PHAS* transcripts) in the RNA-seq data to assess the correlation of phasiRNA and *PHAS* mRNA levels. As many 21-*PHAS* loci are found in large clusters in the rice genome (Johnson et al., 2009) (see Supplementary Table S4), we selected a representative cluster on chromosome 3 for this analysis. Our earlier observations in wild-type rice showed a stage 5 peak for 21-nt phasiRNAs, which reduced slightly at stage 7 (Fig. 3C); in the RNA-seq data at stage 3 and stage 5, 21-*PHAS* transcripts levels were generally consistent with phasiRNA production; however, at stage 7, 21-*PHAS* transcripts levels were very low (Fig. 4) while phasiRNA levels were still high. This is an indication that 21-nt phasiRNAs persist longer than their precursor transcripts, possibly for a prolonged role in later stages of spikelet development. As for *msp1* and *ostd11a* mutants, levels of neither 21-nt phasiRNAs nor 21-*PHAS* transcripts were impacted (stage 5 for both mutants is shown in Supplementary Fig. S3; stage 7 data are not shown but were not appreciably different from the wild-type). Consistent with the precursor molecules consisting of a polyadenylated mRNA paired with an RDR6-derived antisense strand, the strand-specific RNA-seq reads were mapped to only one strand at a given *PHAS* locus (Fig. 4). An examination of the RNA-seq data for 24-*PHAS* loci showed a strong boost in transcript levels from stage 5 to stage 7 in wild-type rice anthers (Supplementary Fig. S4A). However, unlike 21-*PHAS* loci, both 24-nt phasiRNAs and 24-*PHAS* mRNAs were absent in the RNA-seq data from *msp1* and *ostd11a* mutants (Supplementary Fig. S4A, B). In summary, sequencing data revealed the developmental modulation of phasiRNA biogenesis in rice anthers, with 24-phasiRNAs and their precursors disrupted in the *msp1* and *ostd11a* mutants, reflecting an intimate association of these transcripts with anther development and microsporogenesis in rice.

#### *Identification of specialized Argonautes that load reproductive phasiRNAs*


Argonautes (AGOs) are core effector proteins in small RNA-mediated silencing pathways. Different small RNAs are preferentially recruited into specific AGOs, mainly determined by the 5'-terminal nucleotide of the small RNA (Mi et al., 2008). The ten AGO proteins of Arabidopsis are reasonably well studied, but the functional roles of the ~17 to 19 AGOs in grass genomes are less well described (Zhang et al., 2015). The naming for OsAGOs in this study is consistent with the description by Zhang et al. (2015). Moreover, it is still unclear which AGOs recruit the abundant reproductive phasiRNAs of grasses. A recent study in rice showed that the germline-specific AGO protein MEL1 associates with 21-nt phasiRNAs that have 5'-terminal cytosine (Komiya et al., 2014). In addition, the AGO(s) that recruits 24-nt phasiRNAs is still

unknown, although *ZmAGO18b* is enriched in tapetum and germ cells in maize anthers (Zhai et al., 2014), where 24-nt phasiRNAs accumulate (Zhai et al., 2015). Considering the dramatic changes of phasiRNA abundances across different stages of rice spikelet development, we hypothesized that AGO proteins that recruit phasiRNAs may show a gene expression pattern correlated with phasiRNA abundances. Therefore, we examined the expression of all AGOs in rice from stages 3 to 7 (Fig. 5A). We found that *MEL1* peaked at stage 5, the same stage as the peak of accumulation of 21-nt phasiRNAs; therefore, *MEL1* expression was indeed correlated with phasiRNA production. In the RNA-seq data, *MEL1* levels were slightly higher in both mutants (see Supplementary Fig. S5), perhaps because of the excessive number of sporocytes, where *MEL1* is expressed, in mutants (Nonomura et al., 2007). Intriguingly, we found that *OsAGO1d* showed the same pattern as *MEL1*, suggesting a possible functional connection between the OsAGO1d protein and reproductive phasiRNAs (Fig. 5A). Similar to *MEL1*, *OsAGO1d* expression was barely impacted in *msp1* and *ostd11a* mutants (Supplementary Fig. S5).

We speculated that AGOs that accumulate in stage 7 could function with 24-nt phasiRNAs. Gene expression analysis showed that only three *OsAGOs*, namely *OsAGO2b*, *OsAGO5b*, and *OsAGO18*, displayed a substantial up-regulation at stage 7 compared to stages 3 and 5 (Fig. 5B). Since 24-nt phasiRNA accumulation was deficient in *msp1* and *ostd11a* mutants, the result of defects in cell layers important for their biogenesis, it is possible that the *OsAGOs* that load 24-nt phasiRNAs are similarly impacted in both mutants. We found that the levels of *OsAGO2b* and *OsAGO18*, but not *OsAGO5b*, were partially reduced in the *msp1* and *ostd11a* mutants (Fig. 5B), suggesting that OsAGO2b and OsAGO18 are candidates for roles with 24-nt phasiRNAs. To further confirm their specificity to anthers, we checked their expression patterns using published rice anther microarray datasets. Consistent with our RNA-seq data, *OsAGO2b* and *OsAGO18*, but not *OsAGO5b*, showed a meiosis-specific expression pattern in rice anthers (see Supplementary Fig. S6).

#### *Temporal-spatial expression of phasiRNA-associated AGOs revealed by in situ hybridizations*


To connect the temporal specificity of transcript accumulation with spatial patterns, we performed *in situ* hybridizations or examined published images for selected rice *AGO* genes. Published images for *OsAGO2b* demonstrate its accumulation in sporocytes and wall layers of rice anthers at pre-meiotic stages, and in later meiosis stage the transcript is restricted predominantly to the tapetum layer and microspores (Devshwar et al., 2011). We performed *in situ* hybridizations in rice anthers to examine the transcript accumulation patterns of *OsAGO1d* and *OsAGO18* (Fig. 6). These results showed that *OsAGO1d* accumulates highly in the distal epidermis and primary parietal cells of the anther lobe at stage 3. This *OsAGO1d* pattern is reminiscent of miR2118, which accumulates in the distal cells of the epidermis in maize




**Fig. 4.** 21-nt *PHAS* precursor transcripts peak coincidentally with their phasiRNA products. We examined a randomly selected 21-*PHAS* locus on rice chromosome 3 in wild-type (cultivar 9522) rice to assess the peak of abundance relative to the 21-phasiRNAs that peak at stage 5. Because *PHAS* loci are highly clustered in the rice genome, we selected a region of ~24 kb as an example; in this case, the *PHAS* loci are interlaced with repetitive sequences. Each dot is a small RNA; light blue represent 21-nt sRNAs, green represent 22-nt, and orange represent 24-nt. Yellow shaded regions are predicted DNA transposons; pink shaded regions are predicted retrotransposons; orange shaded regions are inverted repeats. The small pink box is an annotated miRNA. The phasiRNA loci are essentially the distinct blocks of 21-nt sRNAs (light blue dots). The strand-specific RNA-seq data is represented as an IGV screenshot; blue bars are top-strand reads, and red bars are bottom-strand reads. There was a paucity of RNA-seq reads from these *PHAS* loci data in stage 7 at this cluster of loci, whereas 21-phasiRNAs were still abundant at stage 7.

anthers (Zhai *et al.*, 2015). In stage 5, *OsAGO1d* accumulates highly in the middle layer and the tapetal layer, which are both differentiated from the primary parietal cells. In stage 7, the level of *OsAGO1d* is much lower than that of stage

5 (Fig. 6A). As for *OsAGO18*, its abundance was enriched in tapetal and sporogenous cells at stages 5 and 7 (Fig. 6B). Therefore, the patterns of *OsAGO1d* and *OsAGO18* across different developmental stages of rice anthers are consistent



**Fig. 5.** Abundance of *OsAGO* transcripts in different stages and backgrounds of anther development. (A) mRNA levels of all nineteen *OsAGO*s in wild-type rice. *OsAGO1d* and *MEL1* are highlighted because only these two *AGO* transcripts peaked at stage 5. (B) *OsAGO2b*, *OsAGO18*, and *OsAGO5b* displayed substantial up-regulation at stage 7 compared to stages 3 and 5. *OsAGO2b* and *OsAGO18*, but not *OsAGO5b*, is defective in the *msp1* and *ostdl1a* mutants at stage 7. Significant differences (Student's *t*-test) are indicated: \**P*<0.01 and \*\**P*<0.05; 'ns' indicates no significant difference.



**Fig. 6.** RNA *in situ* hybridization of *OsAGO1d* and *OsAGO18* in different stages and backgrounds of rice anthers. Expression patterns of *OsAGO1d* (A) and *OsAGO18* (B), performed in stages 3, 5, and 7 of anthers, from either wild-type cultivar 9522 or the *msp1* or *ostdl1a* mutants, as indicated. Each image shows one anther lobe. Cell layers are labeled as in Fig. 1. Scale bars indicate 20  $\mu$ m.



with the results of RNA-seq data from rice spikelets, and these *OsAGO* transcripts are highly correlated both spatially and temporally with reproductive phasiRNAs (comparing our *in situ* with those of the phasiRNAs in maize anthers in [Zhai et al., 2015](#)). Taken together, these analyses suggest a possible functional connection between grass reproductive phasiRNAs and the three AGO proteins encoded by *OsAGO1d*, *OsAGO2b*, and *OsAGO18*.

## Discussion

Like most complex developmental changes, rice spikelet development entails widespread changes in mRNAs and small RNAs. Similar data for mutants in *MSP1* and *OsTDL1A*, which encode interacting proteins with crucial roles in initiating early stage reproductive development, demonstrate the impact of disordered cell specification in rice reproductive tissues ([Yang et al., 2016](#)).

miRNAs play important roles in plant development. We identified miRNAs with differential accumulation patterns during rice spikelet development, including conserved miRNAs (miR156, miR172, and miR164). Among these, the role of miR164 in development is not well studied, although it accumulates in spikelet and floral meristems ([Adam et al., 2011](#)). While variation in miRNA levels in different stages of rice spikelets suggests that miRNAs are active, one interesting observation from the *msp1* and *ostdl1a* mutants is that levels of most miRNAs and their targets were largely not impacted in the mutants, despite the block in development ([Fig. 2B, C; Supplementary Fig. S2](#)). This suggests that these miRNA-involved gene silencing pathways are genetically independent or upstream of the *OsTDL1A-MSP1* pathway.

We also assessed phasiRNA production in rice spikelets. Two classes of phasiRNAs have distinct accumulation patterns in grass anthers ([Zhai et al., 2015](#)); as in maize, we found that rice phasiRNAs peak at specific stages during spikelet development. We showed that miR2118, the trigger of 21-nt phasiRNAs, accumulates to the highest level at stage 5 and drops severely in stage 7, while 21-nt phasiRNAs are relatively slightly retarded, peaking at stage 5 but decreasing only slightly at stage 7. As in maize, miR2275 peaks later (stage 7), at the stage at which 24-nt phasiRNAs reached the highest abundance that we measured. In the *msp1* and *ostdl1a* mutants, 24- but not 21-nt phasiRNAs were depleted – consistent with data from the maize mutant *macl* (the ortholog of rice *OsTDL1A*) ([Wang et al., 2012](#)). Therefore, the timing of phasiRNA biogenesis is conserved in rice and maize, two Poaceae evolutionarily separated by ~50 million years ([Wolfe et al., 1989](#)).

AGO proteins are key catalytic components that associate with small RNAs, and different AGOs function as either RNA binders/slicers or chromatin modifiers by loading different classes of small RNAs ([Waterhouse, 2016](#)). Therefore, to know which AGOs load these reproductive phasiRNAs would greatly help elucidate their functions. Considering that in the mutants 24-nt phasiRNAs were impacted but miRNAs were largely not, we may be able to infer AGOs with roles in phasiRNA function. Compared to 10 AGOs in Arabidopsis,

the rice and maize genomes encode more, 19 and 17 respectively ([Zhang et al., 2015](#)). *AGO5* expression in Arabidopsis is specific to somatic ovule tissues, with a role in megagametogenesis ([Tucker et al., 2012](#)). A potentially conserved function of AGO5 in plant gametogenesis has been shown in rice, as the AGO5 relative MEL1 (*OsAGO5c*) binds 21-nt reproductive phasiRNAs with 5' C ([Komiya et al., 2014](#)). AGO1 in rice has four homologs (*OsAGO1a*, *OsAGO1b*, *OsAGO1c*, and *OsAGO1d*); *OsAGO1a/b/c* predominantly recruit miRNAs and other small RNAs with 5'-terminal uridine ([Wu et al., 2009b](#)). Our RNA-seq data showed that rice *OsAGO1d* accumulates in spikelets, synchronous with *MEL1*, making *OsAGO1d* a strong candidate for further functional analysis. In rice, *OsAGO18* is induced upon viral infection, and has been shown to confer resistance to viruses by sequestering miR168, suppressing *OsAGO1* expression ([Wu et al., 2015](#)). Maize has two homologs of AGO18; *ZmAGO18b* is specific to the tapetum and germ cells ([Zhai et al., 2014](#)). Our data showed that *OsAGO18* and *OsAGO2b* transcripts increase substantially at stage 7, coincident with 24-nt phasiRNA accumulation. Moreover, a recent study on Arabidopsis AGO3 showed that this poorly characterized Argonaute mediates RdDM by binding 24-nt siRNAs ([Zhang et al., 2016](#)). Phylogenetic analysis showed that Arabidopsis AGO3 and rice *OsAGO2b* are close to each other ([Zhang et al., 2015](#)), suggesting that *OsAGO2b* could also recruit 24-nt small RNAs, such as the reproductive phasiRNAs, to regulate epigenetic modifications. Furthermore, *in situ* hybridization results for *OsAGO1d*, *OsAGO18*, and *OsAGO2b* showed that these *OsAGOs* have distinct expression patterns during rice anther development, suggesting that they may associate with either 21- or 24-nt phasiRNAs in a stage-specific manner. In summary, our results suggest a functional relevance between *OsAGO1d*, *OsAGO18*, and *OsAGO2b* and grass reproductive phasiRNAs.

What might be the function of these AGO proteins loaded with phasiRNAs? Prior work on MEL1 (associated with 21-nt phasiRNAs) suggests a role in histone modifications; histone H3 lysine 9 dimethylation (H3K9me2) is decreased in the *mell* mutant ([Nonomura et al., 2007](#)). Yet the reduced H3K9me2 in *mell* could be an indirect effect of other epigenetic changes, as many chromatin modifications in plants and other organisms are intricately linked ([Castel and Martienssen, 2013](#)). Considering the lack of sequence complementarity of phasiRNAs and other regions in the genome ([Zhai et al., 2015](#)), it is possible that reproductive phasiRNAs act primarily in *cis* or impact *cis*-adjacent regions by a spreading mechanism. Indeed, a recent study on maize reported that both 21- and 24-*PHAS* loci showed higher levels of CHH methylation in meiocytes than other tissues, such as seedlings ([Dukowicz-Schulze et al., 2016](#)). This finding suggests that reproductive phasiRNAs may play an important role in chromatin remodeling in *cis* around the stage of meiosis. Overall, the coordinated accumulation of 21- and 24-nt phasiRNAs and several AGO transcripts during rice reproductive development suggests that more detailed and comprehensive analyses of DNA methylation and histone modifications are needed, particularly when coupled with mutants in phasiRNA pathways.



## Supplementary data

Supplementary data are available at *JXB* online.

**Figure S1.** *MSP1* and *OsTDL1A* transcript levels in stage 3 of rice spikelets in the wild-type cultivar 9522 and two mutants.

**Figure S2.** miRNA and target transcript levels in different stages and backgrounds of rice spikelets.

**Figure S3.** 21-nt phasiRNAs and precursor transcripts were unaffected in both *mssl1* and *ostdlla* mutants at stage 5.

**Figure S4.** 24-nt phasiRNAs are strongly impacted in stage 7 spikelets of the two rice mutants.

**Figure S5.** mRNA levels of *MEL1* and *OsAGO1d* in different stages and backgrounds of rice spikelets.

**Figure S6.** Expression of AGOs in different tissues and developmental stages of rice anthers in public microarray datasets.

**Table S1.** Summary information for small RNA and RNA-seq libraries prepared in this study.

**Table S2.** miRNA levels in different libraries.

**Table S3.** Differentially expressed miRNAs in different stages and backgrounds of rice spikelet development.

**Table S4.** 21- and 24-nt phasiRNA abundance from each *PHAS* locus.

## Acknowledgements

Research in the Meyers lab is supported by the US National Science Foundation Plant Genome Research Program (award #1339229). Collaborative research and travel between the Zhang and Meyers lab is supported by the British Council's Global Innovation Initiative, which funds the Global Innovations Network (GIN) (see <https://www.cpiib.ac.uk/GII/> for more details, accessed 23 September 2016). Research in the Zhang lab is supported by National Natural Science Foundation of China (31430009; 31110103915; 32322040; 31271698); China Innovative Research Team, Ministry of Education, and the Program of Introducing Talents of Discipline to Universities (111 Project, B14016); National Key Basic Research Developments Program, Ministry of Science and Technology, China (2013CB126902); the Science and Technology Commission of Shanghai Municipality (grant no. 13JC1408200); Leading Scientist in Agriculture of Shanghai Municipality. We gratefully acknowledge the assistance of Mayumi Nakano with data handling, and Jixian Zhai for helpful discussions.

## References

- Adam H, Marguerettaz M, Qadri R, et al.** 2011. Divergent expression patterns of miR164 and *CUP-SHAPED COTYLEDON* genes in palms and other monocots: implication for the evolution of meristem function in angiosperms. *Molecular Biology & Evolution* **28**, 1439–1454.
- Arikis S, Zhai J, Meyers BC.** 2013. Biogenesis and function of rice small RNAs from non-coding RNA precursors. *Current Opinion in Plant Biology* **16**, 170–179.
- Aukerman MJ, Sakai H.** 2003. Regulation of flowering time and floral organ identity by a MicroRNA and its *APETALA2*-like target genes. *Plant Cell* **15**, 2730–2741.
- Axtell MJ.** 2013. Classification and comparison of small RNAs from plants. *Annual Review of Plant Biology* **64**, 137–159.
- Canales C, Bhatt AM, Scott R, Dickinson H.** 2002. EXS, a putative LRR receptor kinase, regulates male germline cell number and tapetal identity and promotes seed development in Arabidopsis. *Current Biology* **12**, 1718–1727.
- Cao P, Jung KH, Choi D, Hwang D, Zhu J, Ronald PC.** 2012. The Rice Oligonucleotide Array Database: an atlas of rice gene expression. *Rice* **5**, 17.
- Castel SE, Martienssen RA.** 2013. RNA interference in the nucleus: roles for small RNAs in transcription, epigenetics and beyond. *Nature Reviews Genetics* **14**, 100–112.
- Chen X.** 2004. A microRNA as a translational repressor of *APETALA2* in Arabidopsis flower development. *Science* **303**, 2022–2025.
- Chen X.** 2009. Small RNAs and their roles in plant development. *Annual Review of Cellular and Developmental Biology* **25**, 21–44.
- Devshwar P, Bovill WD, Sharma R, Able JA, Kapoor S.** 2011. Analysis of anther transcriptomes to identify genes contributing to meiosis and male gametophyte development in rice. *BMC Plant Biology* **11**, 78.
- Dukowic-Schulze S, Sundararajan A, Ramaraj T, Kianian S, Pawlowski WP, Mudge J, Chen C.** 2016. Novel meiotic miRNAs and indications for a role of phasiRNAs in meiosis. *Frontiers in Plant Science* **7**, 762.
- Fei Q, Xia R, Meyers BC.** 2013. Phased, secondary, small interfering RNAs in posttranscriptional regulatory networks. *The Plant Cell* **25**, 2400–2415.
- Hong L, Tang D, Shen Y, Hu Q, Wang K, Li M, Lu T, Cheng Z.** 2012. *MIL2 (MICROSPOROLESS2)* regulates early cell differentiation in the rice anther. *New Phytologist* **196**, 402–413.
- Jia G, Liu X, Owen HA, Zhao D.** 2008. Signaling of cell fate determination by the TPD1 small protein and EMS1 receptor kinase. *Proceedings of the National Academy of Sciences, USA* **105**, 2220–2225.
- Jiao Y, Wang Y, Xue D, et al.** 2010. Regulation of *OsSPL14* by *OsmiR156* defines ideal plant architecture in rice. *Nature Genetics* **42**, 541–544.
- Johnson C, Kasprzewska A, Tennessen K, Fernandes J, Nan GL, Walbot V, Sundaresan V, Vance V, Bowman LH.** 2009. Clusters and superclusters of phased small RNAs in the developing inflorescence of rice. *Genome Research* **19**, 1429–1440.
- Jones-Rhoades MW, Bartel DP, Bartel B.** 2006. MicroRNAs and their regulatory roles in plants. *Annual Review of Plant Biology* **57**, 19–53.
- Jung KH, Han MJ, Lee YS, Kim YW, Hwang I, Kim MJ, Kim YK, Nahm BH, An G.** 2005. Rice Undeveloped Tapetum1 is a major regulator of early tapetum development. *The Plant Cell* **17**, 2705–2722.
- Komiya R, Ohyanagi H, Niihama M, Watanabe T, Nakano M, Kurata N, Nonomura K.** 2014. Rice germline-specific Argonaute MEL1 protein binds to phasiRNAs generated from more than 700 lincRNAs. *Plant Journal* **78**, 385–397.
- Kouchi H, Hata S.** 1993. Isolation and characterization of novel nodulin cDNAs representing genes expressed at early stages of soybean nodule development. *Molecular & General Genetics* **238**, 106–119.
- Kozomara A, Griffiths-Jones S.** 2014. miRBase: annotating high confidence microRNAs using deep sequencing data. *Nucleic Acids Research* **42**, D68–D73.
- Langmead B, Trapnell C, Pop M, Salzberg SL.** 2009. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biology* **10**, R25.
- Lee DY, An G.** 2012. Two AP2 family genes, *SUPERNUMERARY BRACT (SNB)* and *OsINDETERMINATE SPIKELET 1 (OsIDS1)*, synergistically control inflorescence architecture and floral meristem establishment in rice. *Plant Journal* **69**, 445–461.
- Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, and 1000 Genome Project Data Processing Subgroup.** 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079.
- Li N, Zhang DS, Liu HS, et al.** 2006. The rice tapetum degeneration retardation gene is required for tapetum degradation and anther development. *The Plant Cell* **18**, 2999–3014.
- Li YF, Zheng Y, Addo-Quaye C, Zhang L, Saini A, Jagadeeswaran G, Axtell MJ, Zhang W, Sunkar R.** 2010. Transcriptome-wide identification of microRNA targets in rice. *Plant Journal* **62**, 742–759.
- Luo Y, Guo Z, Li L.** 2013. Evolutionary conservation of microRNA regulatory programs in plant flower development. *Developmental Biology* **380**, 133–144.
- Mi S, Cai T, Hu Y, et al.** 2008. Sorting of small RNAs into Arabidopsis argonaute complexes is directed by the 5' terminal nucleotide. *Cell* **133**, 116–127.
- Miura K, Ikeda M, Matsubara A, Song XJ, Ito M, Asano K, Matsuoka M, Kitano H, Ashikari M.** 2010. *OsSPL14* promotes panicle branching and higher grain productivity in rice. *Nature Genetics* **42**, 545–549.
- Nogueira FT, Madi S, Chitwood DH, Juarez MT, Timmermans MC.** 2007. Two small regulatory RNAs establish opposing fates of a developmental axis. *Genes & Development* **21**, 750–755.

- Nonomura K, Miyoshi K, Eiguchi M, Suzuki T, Miyao A, Hirochika H, Kurata N.** 2003. The *MSP1* gene is necessary to restrict the number of cells entering into male and female sporogenesis and to initiate anther wall formation in rice. *The Plant Cell* **15**, 1728–1739.
- Nonomura K, Morohoshi A, Nakano M, Eiguchi M, Miyao A, Hirochika H, Kurata N.** 2007. A germ cell-specific gene of the *ARGONAUTE* family is essential for the progression of premeiotic mitosis and meiosis during sporogenesis in rice. *The Plant Cell* **19**, 2583–2594.
- Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, Mesirov JP.** 2011. Integrative genomics viewer. *Nature Biotechnology* **29**, 24–26.
- Song X, Li P, Zhai J, et al.** 2012. Roles of DCL4 and DCL3b in rice phased small RNA biogenesis. *Plant Journal* **69**, 462–474.
- Toriba T, Suzaki T, Yamaguchi T, Ohmori Y, Tsukaya H, Hirano HY.** 2010. Distinct regulation of adaxial-abaxial polarity in anther patterning in rice. *The Plant Cell* **22**, 1452–1462.
- Trapnell C, Pachter L, Salzberg SL.** 2009. TopHat: discovering splice junctions with RNA-seq. *Bioinformatics* **25**, 1105–1111.
- Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, Pimentel H, Salzberg SL, Rinn JL, Pachter L.** 2012. Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nature Protocols* **7**, 562–578.
- Tsuji H, Aya K, Ueguchi-Tanaka M, et al.** 2006. GAMYB controls different sets of genes and is differentially regulated by microRNA in aleurone cells and anthers. *Plant Journal* **47**, 427–444.
- Tucker MR, Okada T, Hu Y, Scholefield A, Taylor JM, Koltunow AM.** 2012. Somatic small RNA pathways promote the mitotic events of megagametogenesis during female reproductive development in Arabidopsis. *Development* **139**, 1399–1404.
- Wang CJ, Nan GL, Kelliher T, Timofejeva L, Vernoud V, Golubovskaya IN, Harper L, Egger R, Walbot V, Cande WZ.** 2012. Maize multiple archesporial cells 1 (*mac1*), an ortholog of rice *TDL1A*, modulates cell proliferation and identity in early anther development. *Development* **139**, 2594–2603.
- Waterhouse P.** 2016. RNA interference: A dark horse in the AGO stable. *Nature Plants* **2**, 16059.
- Wolfe KH, Gouy M, Yang YW, Sharp PM, Li WH.** 1989. Date of the monocot–dicot divergence estimated from chloroplast DNA sequence data. *Proceedings of the National Academy of Sciences, USA* **86**, 6201–6205.
- Wu G, Park MY, Conway SR, Wang JW, Weigel D, Poethig RS.** 2009a. The sequential action of miR156 and miR172 regulates developmental timing in Arabidopsis. *Cell* **138**, 750–759.
- Wu J, Yang Z, Wang Y, et al.** 2015. Viral-inducible Argonaute18 confers broad-spectrum virus resistance in rice by sequestering a host microRNA. *eLife* **4**, e05733.
- Wu L, Zhang Q, Zhou H, Ni F, Wu X, Qi Y.** 2009b. Rice microRNA effector complexes and targets. *The Plant Cell* **21**, 3421–3435.
- Xie K, Wu C, Xiong L.** 2006. Genomic organization, differential expression, and interaction of SQUAMOSA promoter-binding-like transcription factors and microRNA156 in rice. *Plant Physiology* **142**, 280–293.
- Yang L, Qian X, Chen M, Fei Q, Meyers BC, Liang W, Zhang D.** 2016. Regulatory role of a receptor-like kinase in specifying anther cell identity. *Plant Physiology* **171**, 2085–2100.
- Yoshida H, Nagato Y.** 2011. Flower development in rice. *Journal of Experimental Botany* **62**, 4719–4730.
- Zhai J, Jeong DH, De Paoli E, et al.** 2011. MicroRNAs as master regulators of the plant *NB-LRR* defense gene family via the production of phased, *trans*-acting siRNAs. *Genes & Development* **25**, 2540–2553.
- Zhai J, Zhang H, Arikrit S, Huang K, Nan GL, Walbot V, Meyers BC.** 2015. Spatiotemporally dynamic, cell-type-dependent premeiotic and meiotic phasiRNAs in maize anthers. *Proceedings of the National Academy of Sciences, USA* **112**, 3146–3151.
- Zhai L, Sun W, Zhang K, Jia H, Liu L, Liu Z, Teng F, Zhang Z.** 2014. Identification and characterization of Argonaute gene family and meiosis-enriched Argonaute during sporogenesis in maize. *Journal of Integrative Plant Biology* **56**, 1042–1052.
- Zhang D, Luo X, Zhu L.** 2011. Cytological analysis and genetic control of rice anther development. *Journal of Genetics & Genomics* **38**, 379–390.
- Zhang D, Wilson ZA.** 2009. Stamen specification and anther development in rice. *Chinese Science Bulletin* **54**, 2342–2353.
- Zhang D, Yang L.** 2014. Specification of tapetum and microsporocyte cells within the anther. *Current Opinion in Plant Biology* **17**, 49–55.
- Zhang H, Xia R, Meyers BC, Walbot V.** 2015. Evolution, functions, and mysteries of plant ARGONAUTE proteins. *Current Opinion in Plant Biology* **27**, 84–90.
- Zhang Z, Liu X, Guo X, Wang X-J, Zhang X.** 2016. Arabidopsis AGO3 predominantly recruits 24-nt small RNAs to regulate epigenetic silencing. *Nature Plants* **2**, 16049.
- Zhao DZ, Wang GF, Speal B, Ma H.** 2002. The *EXCESS MICROSPOROCYTES1* gene encodes a putative leucine-rich repeat receptor protein kinase that controls somatic and reproductive cell fates in the *Arabidopsis* anther. *Genes & Development* **16**, 2021–2031.
- Zhao X, de Palma J, Oane R, Gamuyao R, Luo M, Chaudhury A, Herve P, Xue Q, Bennett J.** 2008. *OsTDL1A* binds to the LRR domain of rice receptor kinase *MSP1*, and is required to limit sporocyte numbers. *Plant Journal* **54**, 375–387.
- Zhou M, Gu L, Li P, Song X, Wei L, Chen Z, Cao X.** 2010. Degradome sequencing reveals endogenous small RNA targets in rice (*Oryza sativa* L. ssp. *indica*). *Frontiers in Biology* **5**, 67–90.
- Zhu QH, Upadhyaya NM, Gubler F, Helliwell CA.** 2009. Over-expression of miR172 causes loss of spikelet determinacy and floral organ abnormalities in rice (*Oryza sativa*). *BMC Plant Biology* **9**, 149.