



Call differentiation in the Limnodynastes tasmaniensis
complex (Anura:Leptodactylidae).

by

J.D.Roberts B.Sc. Honours (Adelaide)

Department of Zoology, University of Adelaide.

A thesis submitted for the degree of Doctor of Philosophy,
November, 1976.

Contents

Summary	vi
Declaration	viii
Acknowledgements	ix
1. The evolution of premating isolating mechanisms: some theories and facts	1
1.1 Introduction	1
1.2 Incidental origin	2
1) The theory	2
2) The evidence supporting the incidental origin hypothesis	3
1.3 Reinforcement	4
1) The theory	4
2) The evidence that reinforcement might produce premating isolating mechanisms	5
a) Simulation and experimental studies	5
b) Reinforcement in natural populations	7
i) Isolation in related sympatric and allopatric species	7
ii) Comparisons of two hybridising species at two points in time	7
iii) Comparisons of premating isolating mechanisms in sympatric and allopatric populations of two species with overlapping distributions	9
iv) Hybrid zones and reinforcement	18
1.4 Differing reproductive environments	20
1) The theory	20
2) The evidence	20
1.5 Discussion	22
1) The theories and the evidence	22
2) The <u>Limnodynastes tasmaniensis</u> complex	24

2.	Call structures in the <u>L.tasmaniensis</u> complex: objective delineation of the call races	25
2.1	Introduction	25
2.2	Methods	26
1)	Call recording and analysis	26
2)	Call components and their measurement	28
a)	Basic call components	28
b)	Derived call components	28
c)	Measurement	29
d)	Temperature correction methods	30
2.3	Results	32
2.4	Discussion	34
1)	The call races	34
2)	Distribution of the call types	35
3.	Call function in <u>L.tasmaniensis</u> : an experimental analysis	37
3.1	Introduction	37
3.2	Methods	39
1)	Experimental series one	39
2)	Experimental series two	42
3)	Effects of observer position	43
3.3	Results	44
1)	Experimental series one	44
2)	Experimental series two	44
a)	Call discrimination trials	44
b)	Ovarian state of responsive females	45
c)	Observer effects on frog movement	45
3.4	Discussion	46
4.	Contacts between the call races of <u>L.tasmaniensis</u>	49
4.1	Introduction	49
4.2	The Coorong contact: the southern and western call races	51
1)	Premating isolating mechanisms	51

	iii
a) Calls	51
b) Breeding and calling seasons	53
i) Field data	53
ii) Laboratory data	55
c) Calling sites used by calling males	56
d) Amplexus position and behaviour in amplexus	57
2) Postmating isolating mechanisms	58
a) Hybridisation and hybrid viability	59
i) Artificial hybridisations	59
ii) Pseudo-natural hybridisations	60
iii) Success and time to metamorphosis	60
b) Behaviour in amplexus and egg mass form	62
3) Geographic distribution and limits on the ranges of the southern and western call races	64
a) Distribution	64
b) Limits on ranges, tadpole transplant experiments	64
c) Limits on ranges, environmental changes	67
4) Discussion	67
4.3 The central Victorian contact: the southern and northern call races	70
1) Geographic distribution	71
2) Call structures	72
3) Artificial hybridisations	73
4) Discussion	73
4.4 The Murray Plains contact: the western and northern call races.	77
1) Premating isolating mechanisms and the nature of the interaction between the western and northern call races	77
a) Call component values and reinforcement	77
b) The interpretation of intermediate call values	80
c) Calls of known hybrids	82
d) Egg size and number of eggs per egg mass	83

	iv
i) Eggs per egg mass	83
ii) Egg size	85
e) Egg mass form and behaviour in amplexus	86
f) Calling sites of males	86
g) Calling and breeding seasons	87
2) Postmating isolating mechanisms	88
3) Geographic distribution, range limiting factors and distribution changes.	91
a) Geographic distribution	91
b) Determinants of range limits for the western and northern call races	93
i) Egg mass transplants	94
ii) Tadpole transplants	94
c) Vegetation, soils and climate	95
d) Salinity as a limiting factor	97
e) Changes in the distribution of the northern call race	99
4) Discussion	102
5. Geographic variation in call structure	111
5.1 Introduction	111
5.2 Variation in the southern call race	113
1) Dominant frequency	113
2) Note duration	117
5.3 Variation in the northern call race	119
1) Dominant frequency	119
2) Pulse repetition rate	120
3) Average notes per call	121
4) Note repetition rate	122
5) Note duration	122
5.4 Variation in the western call race	123
1) Introduction	123

a) Climate	123
b) Vegetation	124
c) Sympatric frog species	125
2) Dominant frequency	126
3) Pulse repetition rate	128
4) Average notes per call	129
5) Note repetition rate	130
6) Note duration	132
5.5 Conclusion	132
6. Appendices	137
7. Bibliography	140

Summary

This thesis analyses call differentiation in the Leptodactylid frog, Limnodynastes tasmaniensis. This analysis assumes that males call to attract females as mates and that calls can therefore also function as premating isolating mechanisms. This was investigated experimentally. Males, females and juveniles could all discriminate between their own and grossly different calls. However, the traditional call discrimination set-up, two loudspeakers with a frog released mid-way, was shown to be inadequate to justify inferences about call function.

Acoustic analysis revealed three call races, western, northern and southern, within the morpho-species, L.tasmaniensis. The races differ most in notes per call, note repetition rate and dominant frequency. The distribution of the three races was mapped. They are largely allopatric but three contact areas exist. The western and northern call races meet in a series of narrow hybrid zones on the Murray plains in South Australia, the northern and southern in a broad intergrade in central Victoria and the southern and western overlap along the Coorong, South Australia.

Evolution of the three races was discussed in terms of three hypotheses. Evidence allegedly supporting these hypotheses was critically discussed in the introduction.

1. Reinforcement: changes in an isolating mechanism result from selection favouring devices which reduce the frequency of interbreeding between two genetically distinct groups whose hybrids are inviable.

Calls and other possible premating isolating mechanisms were analysed in contact areas, and although in all overlaps there was evidence of hybridisation, there was no indication of reinforcement. Hybridisation studies, using combinations of all call races, gave no indication of hybrid inviability suggesting the selective agent required for reinforcement (i.e. gamete wastage) was non-existent. There were no obvious environmental

constraints on the expansion of any of the three contact zones. For the Coorong overlap this was confirmed by the survival of southern tadpoles transplanted to sites outside the normal range of this call race. The possible future of all contacts was discussed.

2. Incidental origin: changes in an isolating mechanism are a correlated response to selection in some other context.

Sampling of call structures in populations from a wide range of environments gave no evidence of incidental origin of call differences between populations within call races.

3. Selective origin: changes in a potential isolating mechanism reflect selection to improve their efficiency as mate attractants in the particular local environment they must function in.

In the northern and western call races, the variance of pulse repetition rate was least in populations with either the greatest number of sympatric anuran species or, where there were synchronic species with call dominant frequencies close to that of L.tasmaniensis. This suggests that acoustic interference may be an important influence on call structure.

The evolution of the three call races was discussed given the above results.

Declaration

This thesis contains no material accepted for the award of any other degree or diploma in this or any other university.

To the best of my knowledge this thesis contains no material previously published or written by another person, except when due reference is made in the text.

22 November, 1976

Acknowledgements

Mike Coates supervised this project, offered advice and criticism during its conduct and laboured through various drafts of the manuscript. Murray Littlejohn suggested I work on the L.tasmaniensis complex. I worked with him in central Victoria. He offered comment or advice on numerous occasions during the conduct of this study and supplied call and distribution data. Mike Tyler introduced me to various areas of anuran biology and was continually available for advice and comment. Michael Bull read and criticised drafts of this thesis. Bob Sharrad alerted me to the presence of L.tasmaniensis in the area west of Morgan and we joined in many successful ventures, both aerial and terrestrial, to the Mount Mary, Morgan region.

The following persons helped in the field and or supplied distribution data. Max King, Dennis King, Rod Wells, Bob Henzell, Phil Withers, Niel Piller, Steve McKillup, the late Michael Smyth, Harry Ehmann, Michael Thompson, Ray Swaby, Brian Cooke, Alan Butler, John Bishop and Kingsley Turner.

C.D.Weiss, K.Smith, S.Eberhard and R.M.Prance kindly allowed me to record frogs and set up experiments on their properties.

To all the above and any who have remained unwittingly anonymous, my sincere thanks. Thanks also to persons in the Zoology Departments of Adelaide and Melbourne Universities (I visited the latter on several occasions) who offered advice or with whom I interacted in either social or scientific settings.

My special thanks to Bev for help in recording calls, in setting out field experiments and for continual encouragement.

I was supported financially by a Commonwealth Post Graduate Research Award.