






Offspring physiology following the use of IVM, IVF and ICSI: a systematic review and meta-analysis of animal studies

Kiri H. Beilby ^{1,*}, Ezra Kneebone¹, Tessa J. Roseboom²,
Indah M. van Marrewijk^{3,4}, Jeremy G. Thompson^{3,5},
Robert J. Norman³, Rebecca L. Robker ³, Ben Willem J. Mol ^{1,6},
and Rui Wang ¹

¹Department of Obstetrics and Gynaecology, Monash University, Melbourne, Australia ²Department of Obstetrics and Gynaecology, Academic Medical Centre, Amsterdam, The Netherlands ³The Robinson Research Institute, School of Biomedicine, University of Adelaide, Adelaide, Australia ⁴Department of Obstetrics and Gynaecology, Erasmus MC University Medical Centre, Rotterdam, The Netherlands ⁵ARC Centre of Excellence for Nanoscale BioPhotonics, University of Adelaide, Adelaide, Australia ⁶Aberdeen Centre for Women's Health Research, University of Aberdeen, Aberdeen, UK

*Correspondence address. Department of Obstetrics and Gynaecology, Monash University, Level 5, Monash Medical Centre, 246 Clayton Road, Clayton, VIC 3168, Australia. E-mail: kiri.beilby@monash.edu  <https://orcid.org/0000-0002-1378-5586>

Submitted on March 29, 2022; resubmitted on November 24, 2022; editorial decision on December 5, 2022


TABLE OF CONTENTS

- Introduction
- Methods
 - Search strategy
 - Eligibility criteria
 - Experimental ART protocols
 - Outcomes
 - Study selection, data extraction and risk of bias assessment
 - Data synthesis
- Results
 - Search results
 - Study characteristics
 - Risk of bias
 - Birthweight
 - Length of gestation
 - Blood pressure
 - Metabolic outcomes
 - Behaviour outcomes
 - Lifespan
- Discussion
 - Summary of key findings
 - Interpretation

Strengths and limitations

- Conclusions

GRAPHICAL ABSTRACT



Offspring produced through IVM/IVF/ICSI can have different physiology to their *in vivo* conceived counterparts.

BACKGROUND: Since the birth of the first baby using IVF technology in 1978, over 10 million children have been conceived via ART. Although most aspects of ARTs were developed in animal models, the introduction of these technologies into clinical practice was performed without comprehensive assessment of their long-term safety. The monitoring of these technologies over time has revealed differences in the physiology of babies produced using ARTs, yet due to the pathology of those presenting for treatment, it is challenging to separate the cause of infertility from the effect of treatments offered. The use of systematic review and meta-analysis to investigate the impacts of the predominant ART interventions used clinically in human populations on animals produced in healthy fertile populations offers an alternative approach to understanding the long-term safety of reproductive technologies.

OBJECTIVE AND RATIONALE: This systematic review and meta-analysis aimed to examine the evidence available from animal studies on physiological outcomes in the offspring conceived after IVF, IVM or ICSI, compared to *in vivo* fertilization, and to provide an overview on the landscape of research in this area.

SEARCH METHODS: PubMed, Embase and Commonwealth Agricultural Bureaux (CAB) Abstracts were searched for relevant studies published until 27 August 2021. Search terms relating to assisted reproductive technology, postnatal outcomes and mammalian animal models were used. Studies that compared postnatal outcomes between *in vitro*-conceived (IVF, ICSI or IVM) and *in vivo*-conceived mammalian animal models were included. *In vivo* conception included mating, artificial insemination, or either of these followed by embryo transfer to a recipient animal with or without *in vitro* culture. Outcomes included birth weight, gestation length, cardiovascular, metabolic and behavioural characteristics and lifespan.

OUTCOMES: A total of 61 studies in five different species (bovine, equine, murine, ovine and non-human primate) met the inclusion criteria. The bovine model was the most frequently used in IVM studies (32/40), while the murine model was mostly used in IVF (17/20) and ICSI (6/8) investigations. Despite considerable heterogeneity, these studies suggest that the use of IVF or maturation results in offspring with higher birthweights and a longer length of gestation, with most of this evidence coming from studies in cattle. These techniques

may also impair glucose and lipid metabolism in male mice. The findings on cardiovascular outcomes and behaviour outcomes were inconsistent across studies.

WIDER IMPLICATIONS: Conception via *in vitro* or *in vivo* means appears to have an influence on measurable outcomes of offspring physiology, manifesting differently across the species studied. Importantly, it can be noted that these measurable differences are noticeable in healthy, fertile animal populations. Thus, common ART interventions may have long-term consequences for those conceived through these techniques, regardless of the pathology underpinning diagnosed infertility. However, due to heterogeneous methods, results and measured outcomes, highlighted in this review, it is difficult to draw firm conclusions. Optimizing animal and human studies that investigate the safety of new reproductive technologies will provide insight into safeguarding the introduction of novel interventions into the clinical setting. Cautiously prescribing the use of ARTs clinically may also be considered to reduce the chance of promoting adverse outcomes in children conceived before long-term safety is confidently documented.

Key words: ICSI / IVM / IVF / ART / assisted reproduction / postnatal outcomes / animal model / systematic review

Introduction

Since its inception, the use of reproductive technologies has resulted in the birth of over 10 million children (ESHRE, 2022). Initially developed to overcome the physical obstacle of blocked fallopian tubes, the application of IVF has now expanded to be used for idiopathic infertility, age-related infertility and male factor infertility as well as fertility preservation. The development of ICSI to further address severe male-factor infertility was introduced clinically in 1992 (Palermo et al., 1992) and widely adopted into practice with little investigation into any potential long-term health effects on offspring. Further, while established for animal application, IVM is increasingly emerging as a technique for clinical use in human infertility, albeit with some variation in protocol compared to animal models where the oocyte is often hormonally 'primed' (treated with exogenous hormones prior to oocyte collection) *in vivo* prior to IVM (Krisher, 2022). Currently, IVF and ICSI are regarded as a set of relatively safe clinical and laboratory procedures that are standardized and available in most countries, with IVM becoming more accessible through practising clinics globally.

The short- and long-term health outcomes of children conceived using IVF and ICSI continue to stimulate research (Painter and Roseboom, 2007; Hart and Norman, 2013; Kamphuis et al., 2014; Pinborg, 2019). This is in part because the Developmental Origins of Health and Diseases (DOHaD) paradigm indicates that there are critical periods of development, including the periconception period where epigenetic reprogramming is taking place, that when perturbed, predispose an adult to cardiovascular and metabolic disorders (Wadhwa et al., 2009). A greater body of evidence is now emerging that suggests altered conditions during periconception can not only influence gamete maturation and preimplantation development but also affect foetal and postnatal growth, as well as adult glucose metabolism, fat deposition and vascular function (Feuer and Rinaudo, 2016).

Existing epidemiological studies, although limited, have shown that children born to infertile parents conceived through ART may be associated with a lower birth weight, higher blood pressure and higher fasting glucose when compared to children conceived spontaneously (Ceelen et al., 2008; Hart and Norman, 2013; Berntsen et al., 2019; Cui et al., 2020). Birth defects and other perinatal outcomes are reported to be more common in ART versus non-ART-conceived children (Pandey et al., 2012; Hansen et al., 2013). Although evidence on cognitive development following ART is subject to methodological limitations, it has been shown that ART, especially ICSI, may be a determinant of child developmental outcomes (Rumbold et al., 2017). In

addition, evidence has been presented to indicate that the incidence of autism spectrum disorder is increased in children conceived by ART (Liu et al., 2017).

These clinical epidemiological studies on outcomes following ART have a major challenge in differentiating 'the chicken and the egg'; it is difficult to attribute the effects on the outcomes to ART or infertility *per se* (Berntsen et al., 2019). Such a challenge cannot be solved by the clinical 'gold standard' method, a randomized controlled trial, as it is impossible to randomize couples with infertility to the natural conception (NC) group due to biological limitations, or to randomize couples without infertility to the ART group due to ethical considerations. Therefore, animal studies play a unique role in this dilemma and animal models are widely used to investigate the impact of specific ART processes on embryo development and postnatal physiology. ART has been used extensively in a wide range of species, not only for research purposes but also for the management of genetics in commercially important agricultural species, and for wildlife conservation. The major advantage of studying ART in animal models is that, unlike in human clinical studies, animals with presumably normal fertility are used, removing the confounding factor of sub-fertility in the population utilizing ART. Furthermore, animal populations are more genetically homogeneous compared to human sub-populations, which may be of benefit in recognizing the true magnitude of variability in treatment outcomes.

Systematic reviews are commonly performed in clinical reproductive medicine, but less common in animal research despite their value to inform pre-clinical, translational and clinical research and practice (Ritskes-Hoitinga et al., 2014). This systematic review aimed to assess the scientific literature reporting animal studies that measured postnatal physiological outcomes following *in vitro* conception (IVF, IVM or ICSI) compared to *in vivo* fertilization and to provide an overview on the landscape of research in this area.

Methods

This review was registered in Prospero: CRD42020191346

Search strategy

Literature reporting on the impact of the mode of conception on offspring health was searched, only to include animal models. We identified studies by searching the PubMed, Embase and *Commonwealth Agricultural Bureaux* (CAB) Abstracts databases from their inception

until 27 August 2021. We used a broad combination of search terms relating to assisted reproductive technology, mammalian animal models and postnatal outcomes (Supplementary Table S1). Only full-text studies in English were included.

Eligibility criteria

To be eligible, studies had to meet all of the following inclusion criteria: (i) conducted in a mammalian animal model; (ii) included animals conceived using an ART intervention, i.e. *in vitro* conception via IVF or ICSI with or without IVM; (iii) included animals conceived following an control group involving an *in vivo* fertilization process: mating/NC, artificial insemination (AI), or NC or AI followed by embryo transfer with or without *in vitro* culture (*in vivo*-ET); and (iv) reported on at least one outcome of interest.

The schematic diagram on the definitions of the ART interventions and *in vivo* controls is presented in Supplementary Fig. S1. Studies reporting on transgenic animals and studies that quantified purely genomic or proteomic measures from organ and tissue samples were excluded.

Experimental ART protocols

Experimental protocols used in the creation of an *in vitro* embryo combine a multitude of steps that vary greatly from the source of the gametes used, to the composition of the culture media chosen, to the micromanipulations applied by the operator. However, all introduce an exposure to IVF, ICSI or IVM. In Supplementary Fig. S1, a comparative summary of the protocols compared in this review is presented.

Outcomes

The outcomes of interest were birth weight, length of gestation, cardiovascular (primarily blood pressure), metabolic (fasting glucose, fasting insulin, area under the curve of glucose and insulin and other metabolites such as lipids) and behavioural measures, and lifespan.

Study selection, data extraction and risk of bias assessment

Potentially eligible articles were first identified by screening titles and abstracts of the search results, followed by a full-text review in Covidence, with both processes conducted independently by at least two authors (K.H.B., E.K., I.M.v.M., J.G.T. and R.L.R.). Differing assessments were resolved by involving another author (K.H.B. or R.W.).

For eligible studies, the following information was extracted by two authors (I.M.v.M., E.K., J.G.T., B.W.J.M. and R.W.) independently: name of the first author, year of publication, country, species, sex, sample size and details on the interventions, controls and outcomes. For all outcome measures, information on method, unit and age of measurements were all extracted. For continuous variables, means and SDs were extracted. If SEs were reported, they were converted to SDs according to a standard approach (Higgins *et al.*, 2022). If the outcomes were only displayed graphically, they were extracted in an online tool, WebPlotDigitizer (version 4.5) (Rohatgi, 2021).

The risk of bias was assessed using a modified SYRACLE risk of bias tool for animal studies (Hooijmans *et al.*, 2014). The following eight domains were evaluated: baseline characteristics, random housing, blinding (performance bias), random outcome assessment, blinding

(detection bias), incomplete outcome data, selective outcome reporting and other sources of bias. We did not assess sequence generation and allocation concealment as randomization may not always be applicable for studies evaluating offspring outcomes.

Two authors (from K.H.B., E.K., T.J.R. and R.W.) independently assessed each included article and discrepancies were solved by consulting another author (K.H.B. or R.W.). The results were graphically displayed using the *robvis* tool (McGuinness and Higgins, 2021).

Data synthesis

To provide an overview on the landscape of research in this area, we plotted an alluvial diagram of included studies by visually linking three categories (year of publication, intervention of interest and species investigated). For continuous outcomes, the summary effects were expressed as mean differences with 95% CIs. Meta-analyses were performed using a random-effects model with the restricted maximum likelihood method. Heterogeneity between studies was quantified using the I^2 statistic. Forest plots were presented for all meta-analyses and different *in vitro* interventions (IVM, IVF and ICSI) were analysed separately when sufficient numbers of studies are available. We explored the source of heterogeneity by stratifying data in subgroup analyses according to different types of controls. We also used a funnel plot to evaluate small-study effects. As *in vivo* fertilization followed by embryo transfer is not practiced in human clinical practice, we performed a sensitivity analysis by limiting the studies to those with an NC or AI control group.

When meta-analyses were impossible due to heterogeneity, we presented forest plots of individual studies without synthesizing the overall estimates. When meta-analyses were impossible due to the missing reporting of variances (SE or SD) in multiple studies, we summarized the means of included studies in Box-and-whisker plots as suggested by the Cochrane handbook (McKenzie and Brennan, 2022). For other circumstances where meta-analyses were impossible, we tabulated the findings of included studies. Results were stratified by the sex of the offspring when possible.

The alluvial diagram was produced in RAWGraphs (Mauri *et al.*, 2017) and data analysed were performed using 'meta' suite in Stata (version 16.1 StataCorpLP, TX, USA).

Results

Search results

The database search returned 5192 articles, of which 61 primary studies met the inclusion criteria after full-text review (Behboodi *et al.*, 1995; Sinclair *et al.*, 1995; Thompson *et al.*, 1995; Holm *et al.*, 1996; Otoi *et al.*, 1996; Kruij and den Daas, 1997; Agca *et al.*, 1998; McEvoy *et al.*, 1998; van Wagtenonk-de Leeuw *et al.*, 1998, 2000; Ptak *et al.*, 1999; Jacobsen *et al.*, 2000a,b, 2002, 2003; Numabe *et al.*, 2000, 2001; Sangild *et al.*, 2000; Yang *et al.*, 2001; Bertolini *et al.*, 2002; Lazzari *et al.*, 2002; Martinez *et al.*, 2002; Ptak *et al.*, 2002; Sakaguchi *et al.*, 2002; Bertolini *et al.*, 2004; Breukelman *et al.*, 2004, 2005; Quaresma *et al.*, 2004; Wolf *et al.*, 2004; Walmsley *et al.*, 2004; Park *et al.*, 2005; Rerat *et al.*, 2005; Givens *et al.*, 2006; Sackett *et al.*, 2006; Hashimoto *et al.*, 2007; Liang *et al.*, 2007; Fernández-Gonzalez *et al.*, 2008; Camargo *et al.*, 2010; Scott *et al.*, 2010; Li *et al.*, 2011;

Kohda et al., 2011; Pimenta-Oliveira et al., 2011; Rexhaj et al., 2013; Wang et al., 2013; Chen et al., 2014a,b; Donjacour et al., 2014; Bonilla et al., 2014; Feuer et al., 2014; Kannampuzha-Francis et al., 2015; Strata et al., 2015; Rexhaj et al., 2015; Cerny et al., 2017; Siqueira et al., 2017; Wang et al., 2017; Valenzuela et al., 2018; Le et al., 2019; Aljahdali et al., 2020; Lewon et al., 2020; Narapareddy et al., 2021; Qin et al., 2021). The PRISMA flow diagram illustrating the process of study screening and selection is shown in Fig. 1.

Study characteristics

The characteristics of the 61 included studies are detailed in Table I and Supplementary Table SII. Briefly, the 61 studies reported on 5 different species (bovine, equine, murine, ovine and non-human primate) published between 1995 and 2021. The research landscape of the animal studies that report on the postnatal outcomes of offspring following the use of ART is presented in Fig. 2. The bovine model was the most frequently used in IVM studies (32/40), while the murine model was mostly chosen for studies that used IVF (17/20) and ICSI (6/8) interventions. There was a shift in the technology studied from IVM in the 1990s to ICSI and IVF from the early 2010s, with the bulk of studies also moving away from bovine and towards murine as a study animal (Fig. 2).

Risk of bias

The complete risk of bias assessment for each included study is presented in Supplementary Fig. S2. While the risk of selection bias was

low in 26/61 studies, it was unclear in the remaining studies where there were baseline differences between the ART and the control group including source of the gametes, breed of the animal models and the timing of the intervention. The animals were housed under the same conditions in 37/61 studies resulting in low risk of bias for that domain, although six studies housed their animals in multiple locations resulting in high risk of bias (Sinclair et al., 1995; Kruij and den Daas, 1997; van Wagtenonk-de Leeuw et al., 1998, 2000; Camargo et al., 2010; Valenzuela et al., 2018). The remaining 18 studies did not report housing conditions. Blinding of caregivers and/or investigators was only reported in 1/61 studies (Siqueira et al., 2017). There was low risk of detection bias amongst the studies due to the objective nature of most of the included outcomes (61/61). The risk of reporting bias was low for most studies (60/61), although it was unclear whether all of one study's prespecified outcomes were reported due to the vague nature of its aim (Otoi et al., 1996). For attrition bias, there was low risk in all but twelve studies (50/61); the risk of bias was unclear in eleven studies as it was not clear if all animals were included in the analysis, and it was high in one study which did not report the reason for attrition (Hashimoto et al. 2007).

Birthweight

Overall, there were 49 studies that reported birthweight (bovine, n = 32; murine, n = 9; ovine, n = 5; equine, n = 1; primate, n = 2). Forest plots depicting our meta-analysis of birthweight between




Figure 1. PRISMA flowchart.

Table 1 Characteristics of included studies.

Study	Country	Species	Intervention	Control
Agca (1998)	USA	Bovine	IVM	AI
Aljahdali (2020)	UK	Murine	IVF	Natural; <i>in vivo</i> -ET
Behboodi (1995)	USA	Bovine	IVM	AI
Bertolini (2002)	USA	Bovine	IVM	<i>In vivo</i> -ET
Bertolini (2004)	USA	Bovine	IVM	<i>In vivo</i> -ET
Bonilla (2014)	USA	Bovine	IVM	AI
Breukelman (2004)	Netherlands	Bovine	IVM	<i>In vivo</i> -ET
Breukelman (2005)	Netherlands	Bovine	IVM	<i>In vivo</i> -ET
Camargo (2010)	Brazil	Bovine	IVM	AI/natural
Cerny (2017)	Switzerland	Murine	IVF	Natural
Chen (2014a)	Australia	Murine	IVF	<i>In vivo</i> -ET
Chen (2014b)	Australia	Murine	IVF	<i>In vivo</i> -ET
Donjacour (2014)	USA	Murine	IVF	Natural
Fernández-Gonzalez (2008)	Spain	Murine	ICSI	<i>In vivo</i> -ET
Feuer (2014)	USA	Murine	IVF	<i>In vivo</i> -ET
Givens (2006)	USA	Bovine	IVM	Natural
Hashimoto (2007)	Japan	Murine	IVF	Natural
Holm (1996)	Australia	Ovine	IVM	<i>In vivo</i> -ET
Jacobsen (2000a)	Denmark	Bovine	IVM	AI
Jacobsen (2000b)	Denmark	Bovine	IVM	AI
Jacobsen (2002)	Denmark	Bovine	IVM	AI
Jacobsen (2003)	Denmark	Bovine	IVM	AI
Kannampuzha-Francis (2015)	USA	Bovine	IVM	AI
Kohda (2011)	Japan	Murine	IVF; ICSI	<i>In vivo</i> -ET
Kruip (1997)	Netherlands	Bovine	IVM	AI
Lazzari (2002)	Italy	Bovine	IVM	AI; <i>in vivo</i> -ET
Le (2019)	China	Murine	IVM; IVF; ICSI	<i>In vivo</i> -ET
Lewon (2020)	USA	Murine	ICSI	Natural
Li (2011)	China	Murine	IVF	Natural
Liang (2007)	China	Bovine	IVM	AI
Martinez (2002)	Argentina	Bovine	IVM	<i>In vivo</i> -ET
McEvoy (1998)	UK	Bovine	IVM	<i>In vivo</i> -ET
Narapareddy (2021)	USA	Murine	IVF	Natural
Numabe (2000)	Japan	Bovine	IVM	<i>In vivo</i> -ET
Numabe (2001)	Japan	Bovine	IVM	<i>In vivo</i> -ET
Otoi (1996)	Japan	Bovine	IVM	AI
Park (2005)	South Korea	Bovine	IVM	<i>In vivo</i> -ET
Pimenta-Oliveira (2011)	Brazil	Bovine	IVM	AI
Ptak (1999)	Italy	Ovine	IVM	<i>In vivo</i> -ET
Ptak (2002)	Italy	Ovine	IVM	Natural; AI
Qin (2021)	China	Murine	IVF	Natural
Quaresma (2004)	Portugal	Bovine	IVM	AI
Rerat (2005)	Switzerland	Bovine	IVM	AI
Rexhaj (2013)	Switzerland	Murine	IVF	AI; natural
Rexhaj (2015)	Switzerland	Murine	IVF	Natural
Sackett (2006)	USA	Primate	IVF; ICSI	AI; natural
Sakaguchi (2002)	Japan	Bovine	IVM	AI
Sangild (2000)	UK	Bovine	IVM	AI

(continued)

Table I Continued

Study	Country	Species	Intervention	Control
Scott (2010)	USA	Murine	IVF; ICSI	Natural
Sinclair (1995)	UK	Bovine	IVM	<i>In vivo</i> -ET
Siqueira (2017)	USA	Bovine	IVF	AI; <i>in vivo</i> -ET
Strata (2015)	USA	Murine	IVF	Natural
Thompson (1995)	New Zealand	Ovine	IVM	Natural
Valenzuela (2018)	USA	Equine	IVM	Natural; <i>in vivo</i> -ET
van Wagtenonk-de Leeuw (1998)	Netherlands	Bovine	IVM	AI
van Wagtenonk-de Leeuw (2000)	Netherlands	Bovine	IVM	AI; <i>in vivo</i> -ET
Walmsley (2004)	Canada	Ovine	IVM	Natural
Wang (2013)	China	Murine	IVF; ICSI	Natural
Wang (2017)	China	Murine	IVM	Natural
Wolf (2004)	USA	Primate	IVF; ICSI	Natural
Yang (2001)	South Korea	Bovine	IVM	AI; <i>in vivo</i> -ET

AI, artificial insemination; ET, embryo transfer.




Figure 2. Alluvial diagram of included studies on the research landscape. The alluvial diagram illustrates animal research on ARTs and postnatal outcomes by visually linking three categories (year of publication, intervention of interest and species investigated). The height of the black rectangle is proportional to the number of studies in each subgroup under each category and the width of curved coloured lines are proportional to the number of studies linking two categories. Note that studies on two or more interventions are considered as separate studies in the diagram and therefore the total number here is 68 instead of 61 (five studies included two interventions and one study included three interventions).




Figure 3. Forest plots for birth weight. (a) Bovine studies. (b) Murine studies. (c) Primate studies.

Continued

different species are presented in Fig. 3a–c and subgroup analyses based on the different types of controls are presented in Supplementary Fig. S3. A funnel plot for small study effects is presented in Supplementary Fig. S4.

The meta-analysis of birthweight in bovine models included 30 studies and showed that birthweights in the IVF and IVM group were both

higher compared to *in vivo* controls (IVF: 1 study, 3810 cattle, mean difference (MD) 0.9 kg, 95% CI 0.25 to 1.55; IVM: 29 studies, 18240 cattle, MD 3.2 kg, 95% CI 2.21 to 4.21, $I^2 = 84\%$; Fig. 3a). The heterogeneity between studies was high but the direction of effect was consistent across studies as shown in Fig. 3a. Subgroup analysis based on types of controls did not reveal differences between




Figure 3. Continued.

subgroups (Supplementary Fig. S3a). The funnel plot was symmetrical overall, indicating no evidence of small-study effects (P for Egger's test = 0.16, Supplementary Fig. S4a). Sensitivity analysis

limiting to NC or AI controls showed similar findings as the main analysis (IVF: 1 study, 3810 cattle, MD 0.9 kg, 95% CI 0.25 to 1.55; IVM: 21 studies, 13018 cattle, MD 3.1 kg, 95% CI 1.8 to 4.4,

$I^2 = 84\%$). Two bovine studies (Sinclair *et al.*, 1995; Kannampuzha-Francis *et al.*, 2015) were not included in our meta-analyses as detailed data were not extractable. They both reported a non-significant difference in birthweight.

Nine studies were included in the meta-analysis of birthweight in murine models. Compared to the control group, ICSI was associated with a small increase in birth weight (1 study, 42 mice, MD 0.37 g, 95% CI 0.29 to 0.46), while the evidence on the difference between IVF and the control group was inconclusive (8 studies, 689 mice, MD 0.12 g, 95% CI -0.03 to 0.27, $I^2 = 96\%$; Fig. 3b). Subgroup analysis showed that IVF was associated with a small or no decrease in birth weight compared to the *in vivo* group (MD -0.10 g, 95% CI -0.22 to 0.02, $I^2 = 87\%$), while IVF was associated with an increase in birth weight compared to the natural group (MD 0.30 g, 95% CI 0.18 to 0.43, $I^2 = 91\%$; P for interaction < 0.001 ; Supplementary Fig. S3b). Although the heterogeneity between studies in the natural subgroup was still high, the direction of effect was consistent across studies. Therefore, the overall high heterogeneity could be partly explained by the choice of controls. When limiting to the NC controls in a sensitivity analysis, both ICSI and IVF were associated with a small increase in birth weight (ICSI: 1 study, 42 mice, MD 0.37 g, 95% CI 0.29 to 0.46; IVF: 4 studies, 400 mice, MD 0.30 g, 95% CI -0.18 to 0.43, $I^2 = 91\%$). In addition, one study (Wang *et al.* 2017), which was not included in meta-analyses because data on variance were not extractable, reported comparable birthweights between IVM and control (natural and *in vivo*-ET).

Two studies reported birth weight in primate models (Wolf *et al.*, 2004; Sackett *et al.*, 2006). Meta-analyses did not show significant differences for ICSI or IVF compared to the control group (2 studies, 109 primates, MD -10.27 g, 95% CI -43.61 to 23.08, $I^2 = 0$; Fig. 3c). Subgroup analysis based on types of controls did not reveal significant differences between subgroups (Supplementary Fig. S3c). As all studies had a natural or AI control group, no sensitivity analysis was performed.

Five studies reported birth weight using an ovine model (Thompson *et al.*, 1995; Holm *et al.*, 1996; Ptak *et al.*, 1999, 2002; Walmsley *et al.*, 2004), and one in an equine model (Valenzuela *et al.*, 2018), with all studies comparing IVM and a control group. Meta-analyses were not performed due to limited extractable data. Four ovine studies reported no significant differences in the birthweight of lambs produced after the use of ART (Thompson *et al.*, 1995; Holm *et al.*, 1996; Ptak *et al.*, 1999, 2002) and one study reported more 'abnormally large lambs' (defined as birth weight more than 99th percentile) in the IVM group, compared to the natural conceived control (Walmsley *et al.*, 2004). No significant differences in birth weight were noted by Valenzuela *et al.* in horses (Valenzuela *et al.*, 2018).

Length of gestation

Overall, 32 studies reported gestational length as an outcome (bovine, $n = 27$; murine, $n = 1$; ovine, $n = 1$; equine, $n = 1$; primate, $n = 2$). A forest plot depicting the meta-analysis is presented in Supplementary Fig. S5 and subgroup analyses based on different types of controls are presented in Supplementary Fig. S6. A funnel plot for small study effects is presented in Supplementary Fig. S4b.

There were 27 studies included in a meta-analysis of bovine data (Supplementary Fig. S5). Overall, our meta-analysis showed no difference in gestational length in the IVF group versus controls (1 study, 3810 cattle, MD 0.20 days, 95% CI -0.47 to 0.84) and a longer gestational length in the IVM group compared to controls (26 studies, 18876 bovine, MD 2.17 days, 95% CI 0.87 to 3.46, $I^2 = 87\%$; Supplementary Fig. S5). Subgroup analysis showed that IVM was associated with an increase in gestational length compared to the AI group (MD 2.71 days, 95% CI 1.27 to 4.16, $I^2 = 83\%$) but was inconclusive about IVM compared to the *in vivo*-ET group (MD 2.31 days, 95% CI -0.03 to 4.66, $I^2 = 93\%$) or the NC group (MD -2.00 days, 95% CI -5.09 to 1.09; Supplementary Fig. S6b). Heterogeneity within subgroups remained high but the direction of effect was consistent in the IVM versus AI subgroup. Therefore, the overall high heterogeneity could be partly explained by the choice of controls. Sensitivity analysis limiting to NC or AI controls showed similar findings as the main analysis (IVF: 1 study, 3810 cattle, MD 0.20 days, 95% CI -0.47 to 0.84; IVM: 19 studies, 13669 bovine, MD 2.33 days, 95% CI 0.93 to 3.72, $I^2 = 84\%$). The funnel plot was symmetrical overall, indicating no evidence of small-study effects (P for Egger's test = 0.95, Supplementary Fig. S4b).

Two studies reported length of gestation in a primate model (Wolf *et al.*, 2004; Sackett *et al.*, 2006), one used an equine model (Valenzuela *et al.*, 2018), one in a murine model (Le *et al.*, 2019) and another used an ovine model (Holm *et al.*, 1996). Meta-analysis of studies in the primate model showed that both IVF and ICSI were associated with a slightly shorter gestational length compared to the AI group (MD -4.28 days, 95% CI -5.63 to -2.93 ; MD -3.00 days, 95% CI -5.15 to -0.85) but the differences were not observed when compared to the NC group (Supplementary Fig. S6b). There was no difference observed between the length of gestation in equine, murine or ovine studies following the use of an ART intervention.

Blood pressure

There were seven included studies that used a mouse model to measure blood pressure as an indicator of cardiovascular health in animals conceived following IVF (including ICSI) and culture and a control group (Table II). Four studies reported the systolic/diastolic and mean blood pressure of offspring at a fixed time point between 9 and 52 weeks of age (Fernández-Gonzalez *et al.*, 2008; Rexhaj *et al.*, 2013; Donjacour *et al.*, 2014), with varied results, as shown in Table II. Two studies recorded arterial blood pressure in 12–14-week-old male mice both at a fixed time point and continuously over a 48-h period using telemetry, and the mean fixed and continuous arterial pressures were significantly higher in mice conceived by IVF (Rexhaj *et al.*, 2013, 2015). One study reported blood pressure in 1.5-year-old mice, with an increased systolic blood pressure in female mice conceived through IVM (Le *et al.*, 2019), but not in male mice or those conceived through IVF or ICSI.

Metabolic outcomes

The outcomes of glucose and lipid metabolism reported in included studies are summarized in Supplementary Table SIII. Glucose metabolism was reported in 16 studies (bovine $n = 3$; murine, $n = 13$) and lipid metabolism was reported in four studies using mouse models.

Table II Studies reporting blood pressure in mouse models.

Study	Intervention	Age	Method of measurement	Outcome (blood pressure)	Findings
Aljahdali (2020)	IVF	9/15/21 weeks	Tail-cuff method	Systolic	↑
Donjacour (2014)	IVF	38 weeks	Tail-cuff method	Systolic	↓
				Diastolic	→
				Mean	→
Fernández-Gonzalez (2008)	ICSI	1 year	Tail-cuff method	Systolic	→
Le (2019)	IVF; ICSI; IVM	1.5 years	Tail-cuff method	Systolic	↑ (IVM, female) → (ICSI; IVF)
				Diastolic	→
				Mean	→
Narapareddy (2021)	IVF	39 weeks	Tail-cuff method	Systolic	→
				Diastolic	→
				Mean	→
Rexhaj (2013)	IVF	12–14 weeks	Telemetry	Mean	↑
Rexhaj (2015)	IVF	12–14 weeks	Telemetry	Mean	↑

All studies used mouse models. ↑ Intervention higher than control; ↓ Intervention lower than control; → no significant differences between groups.

Of the three bovine studies, two measured serum glucose levels within 24 h after birth (Sangild *et al.*, 2000; Bertolini *et al.*, 2002) and one measured glucose and insulin levels on Day 2 after birth (Jacobsen *et al.*, 2000a), all comparing IVM and a control group (Supplementary Table SIII). None of the studies reported significant differences on these outcomes between groups. Among 13 studies reporting outcomes on glucose metabolism in a mouse model, all 13 reported serum fasting glucose levels, 8 reported fasting insulin levels, 11 reported glucose area under the curve (AUC) during a glucose tolerance test and 5 reported insulin AUC. Most studies reporting fasting glucose and insulin did not report information on variance (SD or SE), therefore meta-analysis could not be performed. Instead, a box-and-whisker plot was used to visualize the distribution of effect estimates across studies (Fig. 4a and b). The mean difference on fasting glucose levels between the ART and the control groups varied from -0.7 to 2.4 mmol/l (10 studies, median 0.4) in females and from -0.6 to 2.2 mmol/l (10 studies, median 0.8) in males, with more studies in males showing a higher fasting glucose level in the ART group (Fig. 4a). The mean difference on fasting insulin levels between the ART and the control groups varied from -0.35 to 0.2 ng/ml (median 0.01) in females and from -0.92 to 0.15 ng/ml/l (median -0.04) in males, with more studies in males showing a lower fasting insulin level in the ART group (Fig. 4b).

For glucose and insulin AUC, meta-analyses were not performed due to the variation of age at which the outcomes were measured (8–78 weeks; Supplementary Table SIII). Instead, forest plots without summary estimates were used to visualize the difference in glucose AUC and insulin AUC between groups (Fig. 5a and b).

Eight studies (13 comparisons) reported glucose AUC in female mice, and 8 studies (10 comparisons) reported this outcome in male mice (Fig. 5a). The mean difference for glucose AUC ranged from -200 to 630 mmol/l*2h in studies on females (95% CI of individual studies: -381 to 856) and from -353 to 930 mmol/l*2h in studies on males (95% CI of individual studies: -912 to 1067). More studies

in males showed a higher glucose AUC in the ART group than those in females. Three studies reported data on the insulin AUC following an insulin tolerance test in female mice, and four did the same using male mice (Fig. 5b). The mean difference for insulin AUC ranged from -2.96 to -0.41 ng/ml*2h in studies on females (95% CI of individual studies: -4.27 to 0.49) and from -2.28 to 3.24 ng/ml in studies on males (95% CI of individual studies: -3.44 to 4.49). Studies in female mice showed a lower insulin AUC in the ART group, while studies in male mice showed inconsistent findings.

Four studies reported serum lipid metabolites in mice. Three reported cholesterol and triglycerides levels (Le *et al.*, 2019; Narapareddy *et al.*, 2021; Qin *et al.*, 2021). Meta-analysis was performed in two studies with outcomes measured at 12 and 20 weeks. It showed that IVF did not increase cholesterol levels (MD 0.08, 95% CI -0.24 to 0.41 , $I^2 = 60\%$) compared to naturally conceived group and subgroup analysis based on sex showed that the difference mainly occurred in females (males: MD -0.07 , 95% CI -0.36 to 0.22 , $I^2 = 0$; females: MD 0.30, 95% CI 0.10 to 0.50; P for interaction 0.04; Supplementary Fig. S7a). There was insufficient evidence between IVF and naturally conceived group on triglycerides levels (MD 19.21, 95% CI -0.90 to 39.32 , $I^2 = 66\%$), and subgroup analysis based on sex showed that the difference mainly occurred in males (males: MD 28.60, 95% CI 15.16 to 42.04, $I^2 = 0$; females: MD -2.30 , 95% CI -24.30 to 19.70 ; P for interaction 0.02; Supplementary Fig. S7b). The other study reported data on fatty acid composition in the liver and adipose tissue and found that these outcomes were also altered in male IVF/ICSI mice (Wang *et al.*, 2013).

Behaviour outcomes

Nine studies assessed behavioural characteristics in animals conceived using ART (bovine, $n = 2$; murine, $n = 6$; primate, $n = 1$) (van Wagendonk-de Leeuw *et al.*, 1998; Bertolini *et al.*, 2002; Sackett *et al.*, 2006; Fernández-Gonzalez *et al.*, 2008; Kohda *et al.*, 2011; Li *et al.*, 2011; Strata *et al.*, 2015; Wang *et al.*, 2017; Lewon *et al.*, 2020).




Figure 4. Box-and-whisker plots for fasting glucose and insulin in mice studies. (a) Box-and-whisker plot for fasting glucose levels (mean difference between the intervention and the control, mmol/l). The upper and lower limits of the box, represent the 75th and 25th percentiles, respectively. The line within the box represents the 50th percentile (median), and the whiskers represent the ranges. (b) Box-and-whisker plot for fasting insulin levels (mean difference between the intervention and the control, ng/ml).

There was no difference in the overall cognitive development as observed in rhesus macaque neonates and infants after conception by ICSI, IVF, AI, or NC (Sackett *et al.*, 2006). No difference in spatial learning memory ability between IVF and naturally conceived mice was found at 6–7 weeks of age (Li *et al.*, 2011). Yet, in a complete behavioural analysis of mice, female mice born via *in vivo* conception performed better during acquisition learning than ICSI females, a difference not observed in males, and did not demonstrate the same deficit in implicit memory as ICSI-conceived individuals (Fernández-Gonzalez *et al.*, 2008). In a switching discrimination task and delayed non-matching-to-position memory task, *in vivo* derived males performed better at learning and memory compared to their ICSI produced counterparts, and this was not observed in females (Lewon *et al.*, 2020). While Stata *et al.* (Strata *et al.*, 2015) found that there was no difference in anxiety-like behaviour in mice at ages 10–28 weeks, when additional prenatal and postnatal metabolic stress was added through diet, IVF mice show less anxiety-like behaviour. No difference in newborn behavioural traits such as standing time, suckling time and respiratory distress was observed in IVM compared to *in vivo*-conceived calves (Bertolini *et al.*, 2002). However, the need of a breathing stimulus at birth was significantly increased for calves produced after the use of IVM compared to *in vivo*-conceived animals ($P < 0.05$), especially after a Caesarean delivery. Importantly, this effect only occurred in calves conceived by IVM and using a co-culture cell embryo culture system and not in calves conceived by IVM without this culture system (van Wagtenonk-de Leeuw *et al.*, 2000).

Lifespan

The lifespan of mice conceived after IVF was shorter compared to those conceived naturally, but the difference only found to be statistically significant when they were fed a high fat diet (Rexhaj *et al.*, 2013). Under these conditions, median survival was 582 days ($n = 42$) vs 787 days ($n = 53$) in IVF and *in vivo*-conceived mice, respectively.

Discussion

Summary of key findings

This systematic review and meta-analysis were conducted to identify whether postnatal outcomes in offspring born following *in vitro* conception differed from that of offspring born after *in vivo* conception, as evidenced by animal studies. We identified 61 primary studies that met the inclusion criteria of our search, regarding five species (bovine, murine, ovine, equine and non-human primate). Despite considerable heterogeneity, combined analysis of these studies suggests that the use of ART results in offspring with higher birthweights, and a longer length of gestation, with most of this evidence coming from studies in cattle. Reports on blood pressure were inconsistent, but two of the five studies that measured mean arterial blood pressure demonstrated an increase in Blood pressure (BP) following the use of *in vitro* conception in mice. Studies that measured glucose and insulin sensitivity lacked standardization in both the timing of measurement (age) and the




Figure 5. Forest plots for metabolic outcomes in mice studies. (a) Glucose area under the curve during glucose tolerance test (mmol/l*2 h). (b) Insulin area under the curve during insulin tolerance test (ng/ml*2 h).

methods used, yet indicate sex-specific differences in systemic metabolism after the use of an ART intervention. Serum lipids were also reported to be altered in animals conceived *in vitro*, again in a sex-dependant manner. Aspects of animal behaviour were found to be affected in offspring produced using ART interventions, more specifically ICSI, yet these observations have yet to be replicated between test methods or species. Some of the behavioural characteristics observed also appear to be sex specific. Only one study considered the effect of ART on lifespan, demonstrating this to be shorter for mice conceived after the use of IVF when fed a high-fat diet, compared to spontaneously conceived mice on a high-fat diet.

Interpretation

Assisted reproduction comprises of numerous parallel and in series variations in protocols for clinical and animal systems. ART interventions in our review were limited to IVF, IVM and ICSI and did not include *in vivo* produced embryos followed by embryo transfer with or without *in vitro* culture. This choice was made to mimic technologies commonly used in human clinical practice. Although from a technological perspective, *in vivo* embryos recovered by uterine lavage is completely possible in humans and its use in preimplantation genetic testing - aneuploidy (PGT-A) has been recently reported in a research context (Munné *et al.*, 2020), the underlying ethical challenges are enormously controversial (Lambalk *et al.*, 2020; Oron, 2020; Pennings, 2020).

To make broad comparisons on impact we can at best take these together as a sum of interactions. As shown in both human and animal studies, there appears to be an impact, at least under some experimental designs and conditions, of ART on offspring produced. Our review demonstrates that this impact is taking place regardless of the study cohort/population and can be attributed to the intervention itself. Yet caution must be taken in extrapolating specific results, and here we quite importantly address some of the nuances between clinical and animal studies, areas of caution regarding interpretation, and explanatory mechanisms that may help in understanding the observations made.

The ART used in animal models is overall quite different from clinical approaches in humans. For example, donors and recipients are not the same animal, while in humans, the woman usually plays both roles. Differences in embryo collection methodologies vary greatly for mice, cattle and human populations, as do the hormonal stimulation protocols and culture media additives used, such as growth factors like Epidermal growth factor (EGF) that routinely feature in animal culture systems but are largely unapproved for use in human clinical settings. Similarly, IVM protocols differ between animal and human methodologies where the use of hormonal priming may be used in the case of the latter, making the data hard to extrapolate. The age of animals used should also be highlighted, especially in mouse models where pre-pubertal animals are often used at the time of ovarian stimulation to generate *in vivo* matured oocytes or *in vivo*-derived embryos. In contrast, the use of assisted reproduction in human populations is largely concentrated on older individuals. In mouse studies it is important to control for litter size, as this influences birth weight. Most studies controlled for litter size by matching the number of embryos transferred to pseudopregnant recipients, some considering the *in vivo*-conceived control group with embryo transfer as more suitable than a naturally

conceived group, as the former takes superovulation, litter size and embryo transfer procedure into account. One study removed an oviduct from mice in the control group before mating with a male (Scott *et al.*, 2010), but still litter size ranged from 2 to 10 in naturally conceived mice compared with IVF or ICSI-conceived mice that had litter sizes ranging from 1 to 4. In cattle and sheep studies, many of those reported here used serum or co-culture during *in vitro* culture which itself has been associated with alterations in birth weight (Walker *et al.*, 1992; van Wagendonk-de Leeuw *et al.*, 1998; Hasler, 2000). Yet despite these differences, animal models allow for a focus on specific mechanisms, including epigenetic regulation, to be more closely investigated, and for outcomes with broad variability, such as behaviour to be more easily documented.

Our meta-analysis supports the observation of higher calf birthweight and longer length of gestation when embryos are *in vitro* produced. This phenomenon is in part attributed to the co-culture of these embryos with epithelial cells or serum as a protein source in media, as indicated in a recent review (Duranthon and Chavatte-Palmer, 2018). This practice, while increasing the *in vitro* embryonic yield, alters embryo quality resulting in overgrowth syndrome, as recently reviewed in bovine and ovine studies (Li *et al.*, 2019). A protein source provides important support to *in vitro* culture systems as they prevent the sticking of cells to glass and plastic surfaces which cause mechanical damage, yet even well-defined protein sources such as Bovine serum albumin (BSA) have been shown to influence birthweight (Lazzari *et al.*, 2002), demonstrating a complex scenario regarding balancing all physiological needs. To that point, the issue of difference in oxygen concentration between protocols (5% or 20%) need also be mentioned as an important and common variable, although one not focused on in this review.

Altered fetoplacental development in cattle that are *in vitro* conceived has been linked to placental compensation from impaired vascular development in the early stages of pregnancy, and not directly attributed to length of gestation (Miles *et al.*, 2005). The mechanisms of this observation still require elucidation, however, disruption to the epigenetic processes that occur during embryogenesis appear to contribute to impaired trophoblast differentiation and placentation, leading to disease-related outcomes. This is not just the case as seen in animal studies, but also using human data sets that consider epigenetic changes in candidate genes and genome-wide epigenetic changes in cord blood/placenta and early pregnancy tissues (Mani *et al.*, 2020). Here, the authors examined the link between commonly used ART interventions, changes in DNA methylation and the development of short- and long-term adverse outcomes, and much like the current study concluded that inconsistencies between experimental design and endpoints require further investigation to delineate causal associations.

Foetal growth and birthweight are also changed in murine offspring following the use of ART interventions (Vrooman and Bartolomei, 2017), but not with the same consistent trend as seen in cattle, as supported by our analysis. Rather, there appears to be an intersectional effect of culture system, most commonly in association with culture media composition, but also inbred or outbred animal strains and/or stage at embryo transfer, and this has also been seen to have a sex-specific effect (Bloise *et al.*, 2014; Donjacour *et al.*, 2014; Feuer *et al.*, 2014; Ozil *et al.*, 2017). This acts as a reminder regarding the complexity of the preimplantation environment, including any ARTs

used, and the impact of epigenetic modifications during the sensitive pre- and post-implantation process.

Few studies in cattle measured glucose metabolism ($n=3$) and none found differences between control and treatment groups. A more extensive body of literature exists for glucose metabolism in mice ($n=15$), however, the studies varied in outcome, with a sex-specific effect inconsistently identified. Most notably, male mice produced using ART tended to have higher fasting glucose and lower fasting insulin, and a higher glucose AUC than control animals. Also observed more notably in male offspring was an increase in triglyceride level in ART mice compared to a naturally conceived control group, as shown by meta-analysis in the current study. To identify the molecular mechanisms driving these observed differences, a recent study found 21 genes were differentially regulated in *in vitro* produced mouse embryos, along with a noticeably altered epigenetic signature when compared to their *in vivo*-conceived counterparts. Yet the analysis of male and female embryos separately found minimal differences in epigenetic signature and no gene expression changes (Ruggeri et al., 2020). Given the sexually dimorphic nature of disease prevalence (Ober et al., 2008), the genetic mechanisms that are altered via the use of *in vitro* technologies may manifest differently in male and female phenotypes and may be observable at different times across an animal's lifespan. While the triglyceride levels measured in *in vitro* and *in vivo*-conceived mice differ, the DNA methylation signature in the liver of adult mouse offspring also changes, resulting in hypomethylation of genes involved in metabolism and gene transcription regulation in mice produced via IVF, but no difference between male and female animals (Lira-Albarran et al., 2022). Finally, evidence of the use of media additives, such as melatonin, have been shown to reduce the impact of ART on the glucose metabolism of embryos, demonstrating once again the complex impact of the *in vitro* culture environment (Jia et al., 2022).

Strengths and limitations

The strengths of our systematic review include comprehensive literature search, rigorous data extraction (including data extraction from figures) and a variety of data visualization approaches. There are also some limitations. First, we excluded abstract-only publications due to limited extractable information and five non-English publications. Second, highlighted by the SYRCL tool used, the use of investigator blinding was rarely mentioned and differences in baseline characteristics were only reported in approximately half of the included studies. These might have contributed to performance and selection bias. Third, studies that compared *in vivo* fertilization (NC or AI) followed by *in vitro* culture without IVF versus NC or AI were not included, even if referred to as ART in an animal context. The included interventions in our review (IVF, ICSI and IVF) reflect the use of technologies in humans, where offspring created are done so in combination of IVF and culture. Finally, the use of *in vivo* controls was not commonly reported in studies that observed offspring outcomes following cryopreservation and as such this was not a focus of our review.

Conclusions

There are noticeable differences in some physiological measures made on offspring born after the use of ART in healthy animal populations.

ART use in human populations continues to increase in cycle numbers each year, globally. Improved efficacy and access of treatment has no doubt played a role in its increased application, and healthy children continue to be born as a result. However, given a growing body of evidence, in animal and human studies that offspring born following the use of ART can display differences in physiology compared to those who are spontaneously conceived, circumspect application and further scientific enquiry are required. Greater consideration of the methods and controls used to specifically investigate aspects of *in vitro* culture and fertilization are needed, especially to aid clearer decision-making regarding the clinical application of these technologies.

Supplementary data

Supplementary data are available at *Human Reproduction Update* online.

Data availability

The data underlying this article will be shared on reasonable request to the corresponding author.

Authors' roles

B.W.J.M. and J.G.T. conceived the study. K.H.B., E.K., T.J.R., I.M.v.M., J.G.T., R.J.N., R.L.R., B.W.J.M. and R.W. contributed to the study design and protocol development. I.M.v.M. and E.K. performed the search. I.M.v.M., E.K., B.W.J.M., J.G.T. and R.W. extracted the data. I.M.v.M. and R.W. performed the analysis. E.K., T.J.R., R.W. and K.H.B. performed the quality assessment. I.M.v.M., J.G.T., B.W.J.M., K.H.B. and R.W. drafted the manuscript. K.H.B., E.K., T.J.R., I.M.v.M., J.G.T., R.J.N., R.L.R., B.W.J.M. and R.W. interpreted the data, critically revised the article, and approved the final version.

Funding

B.W.J.M. is supported by an National Health and Medical Research Council (NHMRC) Investigator grant (GNT1176437). R.L.R. is supported by an NHMRC Senior Research Fellowship (1117975). R.W. is supported by an NHMRC Emerging Leadership Investigator grant (2009767).

Conflict of interest

B.W.J.M. reports consultancy for ObsEva, Merck KGaA, Guerbet, iGeonomix & Merck and travel support from Merck. The other authors declare no conflicts of interest.

References

Agca Y, Monson RL, Northey DL, Peschel DE, Schaefer DM, Rutledge JJ. Normal calves from transfer of biopsied, sexed and virified IVP bovine embryos. *Theriogenology* 1998;**50**:129–145.

- Aljahdali A, Airina R, Velazquez MA, Sheth B, Wallen K, Osmond C, Watkins AJ, Eckert JJ, Smyth NR, Fleming TP. The duration of embryo culture after mouse IVF differentially affects cardiovascular and metabolic health in male offspring. *Hum Reprod* 2020;**35**: 2497–2514.
- Behboodi E, Anderson GB, BonDurant RH, Cargill SL, Kreuzer BR, Medrano JF, Murray JD. Birth of large calves that developed from in vitro-derived bovine embryos. *Theriogenology* 1995;**44**:227–232.
- Berntsen S, Soderstrom-Anttila V, Wennerholm UB, Laivuori H, Loft A, Oldereid NB, Romundstad LB, Bergh C, Pinborg A. The health of children conceived by ART: 'the chicken or the egg'. *Hum Reprod Update* 2019;**25**:137–158.
- Bertolini M, Mason JB, Beam SW, Carneiro GF, Sween ML, Kominek DJ, Moyer AL, Famula TR, Sainz RD, Anderson GB. Morphology and morphometry of in vivo- and in vitro-produced bovine concepti from early pregnancy to term and association with high birth weights. *Theriogenology* 2002;**58**:973–994.
- Bertolini M, Moyer AL, Mason JB, Batchelder CA, Hoffert KA, Bertolini LR, Carneiro GF, Cargill SL, Famula TR, Calvert CC et al. Evidence of increased substrate availability to in vitro-derived bovine fetuses and association with accelerated conceptus growth. *Reproduction* 2004;**128**:341–354.
- Bloise E, Feuer SK, Rinaudo PF. Comparative intrauterine development and placental function of ART concepti: implications for human reproductive medicine and animal breeding. *Hum Reprod Update* 2014;**20**:822–839.
- Bonilla L, Block J, Denicol AC, Hansen PJ. Consequences of transfer of an in vitro-produced embryo for the dam and resultant calf. *J Dairy Sci* 2014;**97**:229–239.
- Breukelman SP, Perenyi Z, Ruigh D, Van Wagtenonk-de Leeuw L, Jonker AM, Vermooij FH, Beckers JCM, Van Der Weijden JF, Vos Plam GC, Dieleman SJ et al. Plasma concentrations of bovine pregnancy-associated glycoprotein (bPAG) do not differ during the first 119 days between ongoing pregnancies derived by transfer of in vivo and in vitro produced embryos. *Theriogenology* 2005;**63**: 1378–1389.
- Breukelman SP, Reinders JM, Jonker FH, de Ruigh L, Kaal LM, van Wagtenonk-de Leeuw AM, Vos PL, Dieleman SJ, Beckers JF, Perényi Z et al. Fetometry and fetal heart rates between Day 35 and 108 in bovine pregnancies resulting from transfer of either MOET, IVP-co-culture or IVP-SOF embryos. *Theriogenology* 2004;**61**:867–882.
- Camargo LS, Freitas C, de Sa WF, de Moraes Ferreira A, Serapiao RV, Viana JH. Gestation length, birth weight and offspring gender ratio of in vitro-produced Gyr (*Bos indicus*) cattle embryos. *Anim Reprod Sci* 2010;**120**:10–15.
- Ceelen M, van Weissenbruch MM, Vermeiden JP, van Leeuwen FE, Delemarre-van de Waal HA. Pubertal development in children and adolescents born after IVF and spontaneous conception. *Hum Reprod* 2008;**23**:2791–2798.
- Cerny D, Sartori C, Rimoldi SF, Meister T, Soria R, Bouillet E, Scherrer U, Rexhaj E. Assisted reproductive technologies predispose to insulin resistance and obesity in male mice challenged with a high-fat diet. *Endocrinology* 2017;**158**:1152–1159.
- Chen M, Wu L, Wu F, Wittert GA, Norman RJ, Robker RL, Heilbronn LK. Impaired glucose metabolism in response to high fat diet in female mice conceived by in vitro fertilization (IVF) or ovarian stimulation alone. *PLoS One* 2014b;**9**:e113155.
- Chen M, Wu L, Zhao J, Wu F, Davies MJ, Wittert GA, Norman RJ, Robker RL, Heilbronn LK. Altered glucose metabolism in mouse and humans conceived by IVF. *Diabetes (New York)* 2014a;**63**: 3189–3198.
- Cui L, Zhou W, Xi B, Ma J, Hu J, Fang M, Hu K, Qin Y, You L, Cao Y et al. Increased risk of metabolic dysfunction in children conceived by assisted reproductive technology. *Diabetologia* 2020;**63**: 2150–2157.
- Donjacour A, Liu X, Lin W, Simbulan R, Rinaudo PF. In vitro fertilization affects growth and glucose metabolism in a sex-specific manner in an outbred mouse model. *Biol Reprod* 2014;**90**:80.
- Duranthon V, Chavatte-Palmer P. Long term effects of ART: what do animals tell us? *Mol Reprod Dev* 2018;**85**:348–368.
- ESHRE. ART Fact Sheet. 2022. <https://www.eshre.eu/Europe/Factsheets-and-infographics>.
- Fernández-Gonzalez R, Moreira PN, Pérez-Crespo M, Sánchez-Martín M, Ramirez MA, Pericuesta E, Bilbao A, Bermejo-Alvarez P, de Dios Hourcade J, de Fonseca FR et al. Long-term effects of mouse intracytoplasmic sperm injection with DNA-fragmented sperm on health and behavior of adult offspring. *Biol Reprod* 2008;**78**:761–772.
- Feuer S, Rinaudo P. From embryos to adults: a DOHaD perspective on in vitro fertilization and other assisted reproductive technologies. *Healthcare (Basel)* 2016;**4**:51.
- Feuer SK, Liu X, Donjacour A, Lin W, Simbulan RK, Giritharan G, Plane LD, Kolahi K, Ameri K, Maltepe E et al. Use of a mouse in vitro fertilization model to understand the developmental origins of health and disease hypothesis. *Endocrinology* 2014;**155**: 1956–1969.
- Givens MD, Stringfellow DA, Riddell KP, Galik PK, Carson RL, Riddell MG, Navarre CB. Normal calves produced after transfer of in vitro fertilized embryos cultured with an antiviral compound. *Theriogenology* 2006;**65**:344–355.
- Hansen M, Kurinczuk JJ, Milne E, de Klerk N, Bower C. Assisted reproductive technology and birth defects: a systematic review and meta-analysis. *Hum Reprod Update* 2013;**19**:330–353.
- Hart R, Norman RJ. The longer-term health outcomes for children born as a result of IVF treatment: Part I—General health outcomes. *Hum Reprod Update* 2013;**19**:232–243.
- Hashimoto H, Arai T, Ohnishi Y, Eto T, Ito M, Hioki K, Suzuki R, Yamauchi T, Ohsugi M, Saito M et al. Phenotypes of IRS-2 deficient mice produced by reproductive technology are stable. *Exp Anim* 2007;**56**:149–154.
- Hasler JF. In vitro culture of bovine embryos in Menezes's B2 medium with or without coculture and serum: the normalcy of pregnancies and calves resulting from transferred embryos. *Anim Reprod Sci* 2000;**60-61**:81–91.
- Higgins J, Li T, Deeks J. Chapter 6: choosing effect measures and computing estimates of effect. In: Higgins J, Thomas J, Chandler J, Cumpston M, Li T, Page, M, Welch, V (eds). *Cochrane Handbook for Systematic Reviews of Interventions Version 6.3 (Updated February 2022)*. Cochrane, 2022. www.training.cochrane.org/handbook (20 July 2022, date last accessed).
- Holm P, Walker SK, Seamark RF. Embryo viability, duration of gestation and birth weight in sheep after transfer of in vitro matured

- and in vitro fertilized zygotes cultured in vitro or in vivo. *J Reprod Fertil* 1996;**107**:175–181.
- Hooijmans CR, Rovers MM, de Vries RB, Leenaars M, Ritskes-Hoitinga M, Langendam MW. SYRCLE's risk of bias tool for animal studies. *BMC Med Res Methodol* 2014;**14**:43.
- Jacobsen H, Holm P, Schmidt M, Avery B, Greve T, Callesen H. No peri- and postnatal effects on calves born after transfer of in vitro produced embryos vitrified by the open pulled straw (OPS) method. *Acta Vet Scand* 2003;**44**:87–95.
- Jacobsen H, Sangild PT, Schmidt M, Holm P, Greve T, Callesen H. Macromolecule absorption and cortisol secretion in newborn calves derived from in vitro produced embryos. *Anim Reprod Sci* 2002;**70**:1–11.
- Jacobsen H, Schmidt M, Holm P, Sangild PT, Vajta G, Greve T, Callesen H. Body dimensions and birth and organ weights of calves derived from in vitro produced embryos cultured with or without serum and oviduct epithelium cells. *Theriogenology* 2000b;**53**:1761–1769.
- Jacobsen H, Schmidt M, Hom P, Sangild PT, Greve T, Callesen H. Ease of calving, blood chemistry, insulin and bovine growth hormone of newborn calves derived from embryos produced in vitro in culture systems with serum and co-culture or with PVA. *Theriogenology* 2000a;**54**:147–158.
- Jia Y, Liu W, Bai D, Zhang Y, Li Y, Liu Y, Yin J, Chen Q, Ye M, Zhao Y et al. Melatonin supplementation in the culture medium rescues impaired glucose metabolism in IVF mice offspring. *J Pineal Res* 2022;**72**:e12778.
- Kamphuis EI, Bhattacharya S, van der Veen F, Mol BW, Templeton A; Evidence Based IVF Group. Are we overusing IVF? *BMJ* 2014;**348**:g252.
- Kannampuzha-Francis J, Denicol AC, Loureiro B, Karun K, Ortega MS, Hansen PJ. Exposure to colony stimulating factor 2 during pre-implantation development increases postnatal growth in cattle. *Mol Reprod Dev* 2015;**82**:892–897.
- Kohda T, Ogonuki N, Inoue K, Furuse T, Kaneda H, Suzuki T, Kaneko-Ishino T, Wakayama T, Wakana S, Ogura A et al. Intracytoplasmic sperm injection induces transcriptome perturbation without any transgenerational effect. *Biochem Biophys Res Commun* 2011;**410**:282–288.
- Krisner RL. Present state and future outlook for the application of in vitro oocyte maturation in human infertility treatment. *Biol Reprod* 2022;**106**:235–242.
- Kruip T, den Daas JHG. In vitro produced and cloned embryos: effects on pregnancy, parturition and offspring. *Theriogenology* 1997;**47**:43–52.
- Lambalk CB, Van Wely M, Kirkegaard K, De Geyter C. Ethics beyond ethics. *Hum Reprod* 2020;**35**:1–2.
- Lazzari G, Wrenzycki C, Herrmann D, Duchi R, Kruip T, Niemann H, Galli C. Cellular and molecular deviations in bovine in vitro-produced embryos are related to the large offspring syndrome. *Biol Reprod* 2002;**67**:767–775.
- Le F, Lou HY, Wang QJ, Wang N, Wang LY, Li LJ, Yang XY, Zhan QT, Lou YY, Jin F. Increased hepatic INSIG-SCAP-SREBP expression is associated with cholesterol metabolism disorder in assisted reproductive technology-conceived aged mice. *Reprod Toxicol* 2019;**84**:9–17.
- Lewon M, Peters C, Peterson M, Hayes L, Wang Y, Zheng H, Wang Z, Yan W. Assessment of operant learning and memory in mice born through ICSI. *Hum Reprod* 2020;**35**:2058–2071.
- Li L, Le F, Wang L-y, Xu X-R, Lou H-y, Zheng Y-M, Sheng J-Z, Huang H-F, Jin F. Normal epigenetic inheritance in mice conceived by in vitro fertilization and embryo transfer. *J Zhejiang Univ Sci B* 2011;**12**:796–804.
- Li Y, Donnelly CG, Rivera RM. Overgrowth Syndrome. *Vet Clin North Am Food Anim Pract* 2019;**35**:265–276.
- Liang X, Zhang X, Yang B, Cheng M, Huang F, Pang C, Qing G, Liao C, Wei S, Senatore EM et al. Pregnancy and calving rates following transfer of in-vitro-produced river and FI (river x swamp) buffalo (*Bubalus bubalis*) embryos in recipients on natural oestrus or synchronised for ovulation. *Reprod Fertil Dev* 2007;**19**:670–676.
- Lira-Albarran S, Liu X, Lee SH, Rinaudo P. DNA methylation profile of liver of mice conceived by in vitro fertilization. *J Dev Orig Health Dis* 2022;**13**:358–366.
- Liu L, Gao J, He X, Cai Y, Wang L, Fan X. Association between assisted reproductive technology and the risk of autism spectrum disorders in the offspring: a meta-analysis. *Sci Rep* 2017;**7**:46207.
- Mani S, Ghosh J, Coutifaris C, Sapienza C, Mainigi M. Epigenetic changes and assisted reproductive technologies. *Epigenetics* 2020;**15**:12–25.
- Martinez AG, Valcarcel A, De Las Heras MA, De Matos DG, Furnus C, Brogliatti G. Vitrification of in vitro produced bovine embryos: In vitro and in vivo evaluations. *Anim Reprod Sci* 2002;**73**:11–21.
- Mauri M, Elli T, Caviglia G, Uboldi G, Azzi M. RAWGraphs: a visualisation platform to create open outputs. In: *Proceedings of the 12th Biannual Conference on Italian SIGCHI Chapter*. Association for Computing Machinery, Cagliari, Italy, 2017, Article 28.
- McEvoy TG, Sinclair KD, Broadbent PJ, Goodhand KL, Robinson JJ. Post-natal growth and development of simmental calves derived from in vivo or in vitro embryos. *Reprod Fertil Dev* 1998;**10**:459–464.
- McGuinness LA, Higgins JPT. Risk-of-bias VISualization (robvis): an R package and Shiny web app for visualizing risk-of-bias assessments. *Res Synth Methods* 2021;**12**:55–61.
- McKenzie J, Brennan S. Chapter 12: synthesizing and presenting findings using other methods. In: Higgins J, Thomas J, Chandler J, Cumpston M, Li T, Page M, Welch V (eds). *Cochrane*, 2022. www.training.cochrane.org/handbook (20 July 2022, date last accessed).
- Miles JR, Farin CE, Rodriguez KF, Alexander JE, Farin PW. Effects of embryo culture on angiogenesis and morphometry of bovine placentas during early gestation. *Biol Reprod* 2005;**73**:663–671.
- Munné S, Nakajima ST, Najmabadi S, Sauer MV, Angle MJ, Rivas JL, Mendieta LV, Macaso TM, Sawarkar S, Nadal A et al. First PGT-A using human in vivo blastocysts recovered by uterine lavage: comparison with matched IVF embryo controls. *Hum Reprod* 2020;**35**:70–80.
- Narapareddy L, Rhon-Calderon EA, Vrooman LA, Baeza J, Nguyen DK, Mesaros C, Lan Y, Garcia BA, Schultz RM, Bartolomei MS. Sex-specific effects of in vitro fertilization on adult metabolic outcomes and hepatic transcriptome and proteome in mouse. *FASEB J* 2021;**35**:e21523.

- Numabe T, Oikawa T, Kikuchi T, Horiuchi T. Birth weight and birth rate of heavy calves conceived by transfer of in vitro or in vivo produced bovine embryos. *Anim Reprod Sci* 2000;**64**:13–20.
- Numabe T, Oikawa T, Kikuchi T, Horiuchi T. Birth weight and gestation length of Japanese black calves following transfer of embryos produced in vitro with or without co-culture. *J Vet Med Sci* 2001;**63**:515–519.
- Ober C, Loisel DA, Gilad Y. Sex-specific genetic architecture of human disease. *Nat Rev Genet* 2008;**9**:911–922.
- Oron G. How far should we go in the name of science? *Hum Reprod* 2020;**35**:3–4.
- Otoi T, Yamamoto K, Koyama N, Tachikawa S, Suzuki T. A frozen-thawed in vitro-matured bovine oocyte derived calf with normal growth and fertility. *J Vet Med Sci* 1996;**58**:811–813.
- Ozil J-P, Sainte-Beuve T, Banrezes B. $[Mg^{2+}]_o/[Ca^{2+}]_o$ determines Ca^{2+} response at fertilization: tuning of adult phenotype? *Reproduction* 2017;**154**:675–693.
- Painter RC, Roseboom TJ. Cardiovascular health among children born after assisted reproduction. *Eur J Obstet Gynecol Reprod Biol* 2007;**131**:107–108; author reply 109.
- Palermo G, Joris H, Devroey P, Van Steirteghem AC. Pregnancies after intracytoplasmic injection of single spermatozoon into an oocyte. *Lancet* 1992;**340**:17–18.
- Pandey S, Shetty A, Hamilton M, Bhattacharya S, Maheshwari A. Obstetric and perinatal outcomes in singleton pregnancies resulting from IVF/ICSI: a systematic review and meta-analysis. *Hum Reprod Update* 2012;**18**:485–503.
- Park Y, Kim S, Kim J, Park H, Byun M. The effects of duration of in vitro maturation of bovine oocytes on subsequent development, quality and transfer of embryos. *Theriogenology* 2005;**64**:123–134.
- Pennings G. Uterine lavage: ethics of research and clinical applications. *Hum Reprod* 2020;**35**:1949–1953.
- Pimenta-Oliveira A, Oliveira-Filho JP, Dias A, Goncalves RC. Morbidity-mortality and performance evaluation of Brahman calves from in vitro embryo production. *BMC Vet Res* 2011;**7**:79.
- Pinborg A. Short- and long-term outcomes in children born after assisted reproductive technology. *BJOG* 2019;**126**:145–148.
- Ptak G, Clinton M, Tischner M, Barboni B, Mattioli M, Loi P. Improving delivery and offspring viability of in vitro-produced and cloned sheep embryos. *Biol Reprod* 2002;**67**:1719–1725.
- Ptak G, Loi P, Dattena M, Tischner M, Cappai P. Offspring from one-month-old lambs: studies on the developmental capability of prepubertal oocytes. *Biol Reprod* 1999;**61**:1568–1574.
- Qin N, Zhou Z, Zhao W, Zou K, Shi W, Yu C, Liu X, Dong Z, Mao Y, Liu X et al. Abnormal glucose metabolism in male mice offspring conceived by in vitro fertilization and frozen-thawed embryo transfer. *Front Cell Dev Biol* 2021;**9**:637781.
- Quaresma MA, da Costa LL, Horta AEM, Silva JR. Twinning induction and its effects on embryo-foetal and calf survival, and on reproductive efficiency of Mertolengo cattle kept at pasture. *Rev Port Cienc Vet* 2004;**99**:109–115.
- Rerat M, Zbinden Y, Saner R, Hammon H, Blum JW. In vitro embryo production: Growth performance, feed efficiency, and hematological, metabolic, and endocrine status in calves. *J Dairy Sci* 2005;**88**:2579–2593.
- Rexhaj E, Paoloni-Giacobino A, Rimoldi SF, Fuster DG, Anderegg M, Somm E, Bouillet E, Allemann Y, Sartori C, Scherrer U. Mice generated by in vitro fertilization exhibit vascular dysfunction and shortened life span. *J Clin Invest* 2013;**123**:5052–5060.
- Rexhaj E, Pireva A, Paoloni-Giacobino A, Allemann Y, Cerny D, Dessen P, Sartori C, Scherrer U, Rimoldi SF. Prevention of vascular dysfunction and arterial hypertension in mice generated by assisted reproductive technologies by addition of melatonin to culture media. *Am J Physiol Heart Circ Physiol* 2015;**309**:H1151–H1156.
- Ritskes-Hoitinga M, Leenaars M, Avey M, Rovers M, Scholten R. Systematic reviews of preclinical animal studies can make significant contributions to health care and more transparent translational medicine. *Cochrane Database Syst Rev* 2014;**3**:ED000078.
- Rohatgi A. WebPlotDigitizer. Version 4.5. 2021. <https://automeris.io/WebPlotDigitizer> (15 October 2021, date last accessed).
- Ruggeri E, Lira-Albarran S, Grow EJ, Liu X, Harner R, Maltepe E, Ramalho-Santos M, Donjacour A, Rinaudo P. Sex-specific epigenetic profile of inner cell mass of mice conceived in vivo or by IVF. *Mol Hum Reprod* 2020;**26**:866–878.
- Rumbold AR, Moore VM, Whitrow MJ, Oswald TK, Moran LJ, Fernandez RC, Barnhart KT, Davies MJ. The impact of specific fertility treatments on cognitive development in childhood and adolescence: a systematic review. *Hum Reprod* 2017;**32**:1489–1507.
- Sackett G, Ruppenthal G, Hewitson L, Simerly C, Schatten G. Neonatal behavior and infant cognitive development in rhesus macaques produced by assisted reproductive technologies. *Dev Psychobiol* 2006;**48**:243–265.
- Sakaguchi M, Geshi M, Hamano S, Yonai M, Nagai T. Embryonic and calving losses in bovine mixed-breed twins induced by transfer of in vitro-produced embryos to bred recipients. *Anim Reprod Sci* 2002;**72**:209–221.
- Sangild PT, Schmidt M, Jacobsen H, Fowden AL, Forhead A, Avery B, Greve T. Blood chemistry, nutrient metabolism, and organ weights in fetal and newborn calves derived from in vitro-produced bovine embryos. *Biol Reprod* 2000;**62**:1495–1504.
- Scott KA, Yamazaki Y, Yamamoto M, Lin Y, Melhorn SJ, Krause EG, Woods SC, Yanagimachi R, Sakai RR, Tamashiro KL. Glucose parameters are altered in mouse offspring produced by assisted reproductive technologies and somatic cell nuclear transfer. *Biol Reprod* 2010;**83**:220–227.
- Sinclair KD, Broadbent PJ, Dolman DF, Watt RG, Mullan JS. Establishing twin pregnancies in cattle by embryo transfer. *Anim Sci* 1995;**61**:25–33.
- Siqueira LGB, Dikmen S, Ortega MS, Hansen PJ. Postnatal phenotype of dairy cows is altered by in vitro embryo production using reverse X-sorted semen. *J Dairy Sci* 2017;**100**:5899–5908.
- Strata F, Giritharan G, Sebastiano FD, Piane LD, Kao C-N, Donjacour A, Rinaudo P. Behavior and brain gene expression changes in mice exposed to preimplantation and prenatal stress. *Reprod Sci* 2015;**22**:23–30.
- Thompson JG, Gardner DK, Pugh PA, McMillan WH, Tervit HR. Lamb birth weight is affected by culture system utilized during in vitro pre-elongation development of ovine embryos. *Biol Reprod* 1995;**53**:1385–1391.
- Valenzuela OA, Couturier-Tarrade A, Choi YH, Aubriere MC, Ritthaler J, Chavatte-Palmer P, Hinrichs K. Impact of equine assisted reproductive technologies (standard embryo transfer or intracytoplasmic sperm injection (ICSI) with in vitro culture and

- embryo transfer) on placenta and foal morphometry and placental gene expression. *Reprod Fertil Dev* 2018;**30**:371–379.
- van Wagtenonk-de Leeuw AM, Aerts BJ, den Daas JH. Abnormal offspring following in vitro production of bovine preimplantation embryos: a field study. *Theriogenology* 1998;**49**:883–894.
- van Wagtenonk-de Leeuw AM, Mullaart E, de Roos AP, Merton JS, den Daas JH, Kemp B, de Ruigh L. Effects of different reproduction techniques: AI MOET or IVP, on health and welfare of bovine offspring. *Theriogenology* 2000;**53**:575–597.
- Vrooman LA, Bartolomei MS. Can assisted reproductive technologies cause adult-onset disease? Evidence from human and mouse. *Reprod Toxicol* 2017;**68**:72–84.
- Wadhwa PD, Buss C, Entringer S, Swanson JM. Developmental origins of health and disease: brief history of the approach and current focus on epigenetic mechanisms. *Semin Reprod Med* 2009;**27**:358–368.
- Walker SK, Heard TM, Seamark RF. In vitro culture of sheep embryos without co-culture: successes and perspectives. *Theriogenology* 1992;**37**:111–126.
- Walmsley SE, Buckrell BC, Buschbeck C, Rumph N, Pollard JW. Rate of abnormalities in lambs from in vitro produced embryos transferred on Day 2 compared with Day 6 postfertilization. *Theriogenology* 2004;**62**:195–206.
- Wang L-Y, Le F, Wang N, Li L, Liu X-Z, Zheng Y-M, Lou H-Y, Xu X-R, Chen Y-L, Zhu X-M et al. Alteration of fatty acid metabolism in the liver, adipose tissue, and testis of male mice conceived through assisted reproductive technologies: fatty acid metabolism in ART mice. *Lipids Health Dis* 2013;**12**:5.
- Wang N, Ren CE, Lou YY, Le F, Wang LY, Liu XZ, Zhan QT, Mao LN, Lou HY, Jin F. Inter-generational effects of the in vitro maturation technique on pregnancy outcomes, early development, and cognition of offspring in mouse model. *Clin Chim Acta* 2017;**473**:218–227.
- Wolf DP, Thormahlen S, Ramsey C, Yeoman RR, Fanton J, Mitalipov S. Use of assisted reproductive technologies in the propagation of rhesus macaque offspring I. *Biol Reprod* 2004;**71**:486–493.
- Yang BS, Im GS, Park SJ. Characteristics of Korean native, Hanwoo, calves produced by transfer of in vitro produced embryos. *Anim Reprod Sci* 2001;**67**:153–158.