

PUBLISHED VERSION

Wang, He; Leigh, James.

Effects of nitric oxide synthase inhibitor - Nitro-L-Arginine Methyl Ester, on silica-induced inflammatory reaction and apoptosis, *Particle and Fibre Technology*, 2006; 3:14.

© 2006 Wang and Leigh; licensee BioMed Central Ltd.

PERMISSIONS

<http://www.biomedcentral.com/info/about/license>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

BioMed Central Open Access license agreement

Brief summary of the agreement:

Anyone is free:

- to copy, distribute, and display the work;
- to make derivative works;
- to make commercial use of the work;

Under the following conditions: Attribution

- the original author must be given credit;
- for any reuse or distribution, it must be made clear to others what the license terms of this work are;
- any of these conditions can be waived if the authors gives permission.

2nd May 2011

<http://hdl.handle.net/2440/35653>

Research

Open Access

Effects of nitric oxide synthase inhibitor ω -Nitro-L-Arginine Methyl Ester, on silica-induced inflammatory reaction and apoptosis

He Wang*¹ and James Leigh²

Address: ¹Discipline of Public Health, University of Adelaide, 10 Pulteney Street, Adelaide, 5005 SA, Australia and ²School of Public Health, University of Sydney, Sydney, 2006 NSW, Australia

Email: He Wang* - he.wang@adelaide.edu.au; James Leigh - jleigh@bigpond.com

* Corresponding author

Published: 07 November 2006

Received: 24 December 2005

Particle and Fibre Toxicology 2006, **3**:14 doi:10.1186/1743-8977-3-14

Accepted: 07 November 2006

This article is available from: <http://www.particleandfibretoxicology.com/content/3/1/14>

© 2006 Wang and Leigh; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Although nitric oxide is overproduced by macrophages and neutrophils after exposure to silica, its role in silica-induced inflammatory reaction and apoptosis needs further clarification. In this study, rats were intratracheally instilled with either silica suspension or saline to examine inflammatory reactions and intraperitoneally injected with ω -nitro-L-arginine methyl ester (L-NAME), an inhibitor of nitric oxide synthases, or saline to examine the possible role of nitric oxide production in the reaction.

Results: Results showed that silica instillation induced a strong inflammatory reaction indicated by increased total cell number, number of neutrophils, protein concentration and lactate dehydrogenase (LDH) activity in bronchoalveolar lavage fluid (BALF). There were no significant differences in these indices between silica-instilled groups with and without L-NAME injection ($p > 0.05$) except LDH level. The results also showed that apoptotic leucocytes were identified in BALF cells of silica-instilled groups whereas no significant difference was found between silica-instilled groups with and without L-NAME injection in the apoptotic reaction ($p > 0.05$). Silica instillation significantly increased the level of BALF nitrite/nitrate and L-NAME injection reduced this increase.

Conclusion: Intratracheal instillation of silica caused an obvious inflammatory reaction and leucocyte apoptosis, but these reactions were not influenced by intraperitoneal injection of L-NAME and reduced production of NO. This supports the possibility that silica-induced lung inflammation and BALF cell apoptosis are via NO-independent mechanisms.

Background

Silica exposure results in an initial inflammatory reaction and subsequent fibrosis. During this process, various agents such as cytokines and free radicals are produced and these agents in turn regulate the development of the inflammation and fibrosis [1]. Nitric oxide (NO), a small molecule with multiple biologic functions, has been shown to be overproduced by alveolar macrophages and neutrophils as well as other cell types after exposure to

intratracheal instillation of silica [2-4]. It has also been demonstrated that γ -interferon and TNF- α , which are produced in silica-induced response [5], can induce synthesis of NO [6].

NO is a molecule that can readily pass through the cell membrane and exert its action on cells. It is involved in vessel dilatation, inhibition of platelet aggregation [7] and host defence. It is also an apoptosis inducer for some cell

types [8]. Since silica-induced apoptosis of bronchoalveolar leucocytes has already been demonstrated [9] and probably has a role in the evolution of silica-induced inflammation and fibrosis, NO may induce leucocyte apoptosis to regulate these pathological reactions. However, there are also evidences that some apoptotic change occurs via a nitric oxide-independent pathway [10,11]. It is not known whether leucocyte apoptosis in silica-induced inflammation occurs via the nitric oxide-dependent or independent pathway.

Although NO seems to be beneficial by induction of apoptosis in inflammatory cells, the NO produced in a silica-induced lung reaction may be harmful. That is because nitric oxide can combine with superoxide to form peroxynitrite free radical and this substance can cause severe lung damage [2,12,13]. Indeed, other harmful agents such as asbestos fibres [14] and ozone [15], which induce lung damage, have been demonstrated to enhance production of NO in vitro and in vivo although the role of NO in the process is uncertain. Even in the case of peroxynitrite, its exact role in pathological processes is contradictive. Current knowledge indicates that both NO and peroxynitrite may have dual or beneficial and harmful effects in inflammatory reactions depending on situation.

ω -nitro-L-arginine methyl ester (L-NAME) is a NO synthase inhibitor and inhibits the production of NO by inducible NO synthase and constitutive NO synthase. In silica-induced inflammatory reaction, inducible NO synthase gene expression increases in alveolar leucocytes and the production of NO is likely to be via the increased gene expression of NO synthase [3]. In the present study, silica-induced inflammation and apoptosis were quantified in intratracheally (it) silica-instilled rats with and without L-NAME injection to see if this compound could inhibit the inflammatory reaction and apoptosis.

Results

Inflammatory reaction

Intratracheal instillation of silica induced an obvious inflammatory reaction in silica-instilled rats but not in the saline-instilled rats. Intraperitoneal injection (ip) of L-NAME in a dosage of 15 mg/kg/day did not influence the inflammatory reaction significantly. There was no significant difference in total cell number of bronchoalveolar lavage fluid (BALF) between (it) silica + (ip) saline and (it) silica + (ip) L-NAME groups ($p > 0.05$, Figure 1). Both silica-instilled groups showed larger numbers of total cells than the saline-instilled groups but the difference was not statistically significant possibly due to the large individual variation in silica-instilled groups.

The number of neutrophils in BALF increased markedly in both of the silica-instilled groups compared with saline-




Figure 1

Total cell number in bronchoalveolar lavage fluid of rats after intratracheal instillation of silica or saline and intraperitoneal injection of saline or ω -nitro-L-arginine methyl ester (L-NAME). QL = silica (quartz) + L-NAME; QS = silica + saline; SL = saline + L-NAME; SS = saline + saline (These abbreviations also apply to subsequent figures) (mean + se).

instilled rats (Figure 2). There were no significant differences between saline- and L-NAME-treated groups in either silica-instilled or saline-instilled rats.

The number of macrophages in silica-instilled rats decreased significantly compared with saline-instilled groups ($p < 0.01$). This might be a reflection of increased death of alveolar macrophages because of silica toxicity. L-NAME had no influence on the number of macrophages in both silica- and saline-instilled groups (Figure 3).

The protein concentration in BALF of silica-instilled groups significantly ($p < 0.05$) increased compared with saline-instilled groups but no significant difference could be detected between the silica-instilled groups with and without L-NAME injection (Figure 4).

The lung weight of silica-instilled groups was significantly ($p < 0.05$) higher than that of saline-instilled groups (Figure 5), but no significant difference was found between the L-NAME and non-L-NAME silica-instilled groups ($p > 0.05$).

The lactate dehydrogenase (LDH) activity in BALF from silica-instilled groups was significantly ($p < 0.05$) higher than that of saline-instilled groups and L-NAME injection reduced the increase of LDH activity induced by silica instillation from 140 U/L to 95 U/L ($p < 0.05$, Figure 6).




Figure 2
Percentage of neutrophils in BALF cells of rats after intratracheal instillation of silica or saline and intraperitoneal injection of saline or ω -nitro-L-arginine methyl ester (L-NAME) (mean + se). * There is a statistically significant difference in comparison with control groups.




Figure 4
Protein concentration in bronchoalveolar lavage fluid of rats after intratracheal instillation of silica or saline and intraperitoneal injection of saline or ω -nitro-L-arginine methyl ester (L-NAME) (mean + se). * There is a statistically significant difference in comparison with control groups.




Figure 3
Percentage of macrophages in BALF cells of rats after intratracheal instillation of silica or saline and intraperitoneal injection of saline or ω -nitro-L-arginine methyl ester (L-NAME) (mean + se). * There is a statistically significant difference in comparison with control groups.





Figure 5
Lung weight of rats after intratracheal instillation of silica or saline and intraperitoneal injection of saline or ω -nitro-L-arginine methyl ester (L-NAME) (mean + se). * There is a statistically significant difference in comparison with control groups.

**Figure 6**

LDH activity in bronchoalveolar lavage fluid of rats after intratracheal instillation of silica or saline and intraperitoneal injection of saline or ω -nitro-L-arginine methyl ester (L-NAME) (mean + se). * There is a statistically significant difference in comparison with control groups. ** There is a statistically significant difference in comparison with saline-injected group.

Apoptotic Reaction


A number of apoptotic leucocytes were identified in BALF cells of both silica-instilled groups. The apoptotic cells were mainly neutrophils by morphology. Engulfment of apoptotic cells by macrophages was also identified. Comparison of the proportion of apoptotic leucocytes (Figure 7), the percentage of apoptotic neutrophils (Figure 8) and the proportion of macrophages with engulfed apoptotic cells (Figure 9) between the two silica-instilled groups, indicated that the differences were not significant ($p > 0.05$). In saline-instilled groups, apoptotic cells were extremely few.

Nitrite/nitrate level in BALF

Nitrite/nitrate can be detected in BALF from all the experimental animals. Silica instillation increased the level of nitrite/nitrate in BALF significantly two and four weeks after instillation and L-NAME injection reduced the increase by about 40% at two weeks time point and 55% at four weeks time point (Figure 10).

Discussion


In this study, a strong inflammatory reaction was induced following silica instillation. At the administered dosage, L-NAME failed to influence the silica-induced inflammation indices. The L-NAME dosage was above the effective dosage in similar studies for the inhibition of lung injury [16-19] and actually reduced the level of nitrite/nitrate in BALF from silica-instilled rats in this experiment.

**Figure 7**


Proportion of apoptotic leucocytes in BALF leucocytes of rats after intratracheal instillation of silica or saline and intraperitoneal injection of saline or ω -nitro-L-arginine methyl ester (L-NAME) (mean + se). * There is a statistically significant difference in comparison with control groups.

The results in this study may be an indication that although nitric oxide may be an important marker in the development of lung injury, it does not play a pivotal role in silica-induced inflammation under the conditions of this experiment. In a previous study with a rat model [20], intratracheal instillation of silica (100 mg/kg BW) caused overproduction of both superoxide and nitric oxide 24 hours later. Nitric oxide has been demonstrated to be able to combine with superoxide to form peroxynitrite which is a much stronger harmful agent to lung tissue [21,22]. However, NO synthase inhibitor L-NAME did not reduce lung inflammatory reactions in the present study although the level of nitrite/nitrate in BALF from silica-instilled rats was reduced by the drug. L-NAME had also been shown to decrease NO as well as peroxynitrite production in silica-induced acute lung injury [20] in a previous study. The damaging role of NO or peroxynitrite in silica-induced lung injury could not be verified by injection of L-NAME which has been widely used as a NO synthase inhibitor. The dependency of silica-induced lung inflammation on NO or peroxynitrite is open to question.

NO has been demonstrated to cause apoptosis [23]. Induction of inducible NO synthase by exogenous agent can also lead to cell death by apoptosis [24]. Macrophages cause apoptosis of other cell types through the release of nitric oxide [25]. In silica-induced inflammation, increase in inducible nitric oxide synthase activity, nitric oxide overproduction and leucocyte apoptosis have all been demonstrated [26,27]. This suggests that there is a probability that the silica-induced nitric oxide might induce neutrophil apoptosis to regulate the evolution of the sil-

**Figure 8**

Percentage of apoptotic neutrophils in BALF neutrophils of rats after intratracheal instillation of silica or saline and intraperitoneal injection of saline or ω -nitro-L-arginine methyl ester (L-NAME) (mean + se). * There is a statistically significant difference in comparison with control groups.

**Figure 9**

Proportion of pulmonary alveolar macrophages with engulfed apoptotic neutrophils in BALF macrophages of rats after intratracheal instillation of silica or saline and intraperitoneal injection of saline or ω -nitro-L-arginine methyl ester (L-NAME) (mean + se). * There is a statistically significant difference in comparison with control groups.

ica-induced inflammation. However, in this experiment, L-NAME did not influence leucocyte apoptosis significantly. Since the dosage is well above effective and actually reduced the BALF level of nitrite/nitrate, it is possible that nitric oxide plays only a minor role in the leucocyte apoptosis of silica-induced inflammation.


It is noted that chemicals can cause cell apoptosis by either nitric oxide-dependent or nitric oxide-independent mechanisms [28]. For example, bisphosphonates induce apoptosis in mouse macrophage-like cells *in vitro* by an nitric oxide-independent pathway [29]. These workers demonstrated that the apoptosis induced by bisphosphonates was not due to increased synthesis of NO and could not be prevented by inhibitors of NO synthase. NO also failed to induce apoptosis in oligodendrocytes [30].

Peroxynitrite can be produced in silica-induced lung injury and this endogenous substance can induce either necrosis [31,32] or apoptosis[33-36]. In this experiment, L-NAME did not influence either leucocyte apoptosis or lung inflammation significantly. Since peroxynitrite might be overproduced in silica-induced inflammation of this study and L-NAME actually reduced the overproduction of NO, there is a possibility that leucocyte apoptosis in silica-induced inflammation was via a peroxynitrite-independent pathway.


It is interesting that L-NAME at an effective dosage, failed to influence silica-induced inflammation as well as leucocyte apoptosis. This supports the hypothesis that the regu-

lation of apoptosis might influence the evolution of silica-induced inflammation. For example, in a previous study, it was demonstrated that enhancement of apoptosis by IL-10 accelerated the resolution of acute inflammation induced by LPS [37]. It was noted that L-NAME inhibited silica-induced LDH activity increase in BALF in this experiment. This inhibition may also be related to a NO-independent pathway since it has been demonstrated that L-NAME can reduce LDH release by cultured cells independent of NO reduction [38,39].

In regard to the role of NO in injuries caused by various invading agents, conflicting results have been presented. A number of experiments showed that NO was harmful to the exposed cells or tissues [2] whereas some experiments indicated that NO actually played a protective role to the exposed tissues or cells [40,41]. Moreover, in an *in vitro* study NO was demonstrated to play a dichotomous role in asbestos induced harmful effects [42]. The result of this study was the outcome of four weeks after one off exposure of silica and continuing administration of L-NAME. Although this study did not directly measure the activity of NO synthases, the production of NO estimated by nitrite/nitrate level in BALF was dramatically reduced by the injection of L-NAME. Since it has been reviewed that direct link between silica-induced NO production and pathogenesis was obtained using iNOS knockout mice, and iNOS knockout provided protection in sub-chronic exposure of silica but not in acute exposure [43], the complexity of NO synthases in silica-induced inflammation is shown. Further studies are needed to reveal the exact role of NO in silica-induced harmful effects.



(a)



(b)

Figure 10

(a) Concentration of NO₂⁻/NO₃⁻ (µM) in BALF of rats 2 weeks after intratracheal instillation of silica or saline and intraperitoneal injection of saline or ω-nitro-L-arginine methyl ester (L-NAME) (mean + se). ** There is a statistically significant difference in comparison with control groups. * There is a statistically significant difference in comparison with instillation of silica and injection of saline group. (b) Concentration of NO₂⁻/NO₃⁻ (µM) in BALF of rats 4 weeks after intratracheal instillation of silica or saline and intraperitoneal injection of saline or ω-nitro-L-arginine methyl ester (L-NAME) (mean + se). ** There is a statistically significant difference in comparison with control group. * There is a statistically significant difference in comparison with instillation of silica and injection of saline group.

Conclusion

This study demonstrated that L-NAME, a nitric oxide synthase inhibitor, did not influence the inflammatory reaction and apoptosis of neutrophils after exposure to intratracheal silica in rats at the dosage of 15 mg/kg/day for four weeks. Because this dosage was more than 10 times of L-NAME ED 50 and it actually reduced the production of NO in silica-instilled lungs, NO probably played little role in the development of inflammation and leucocyte apoptosis in silica-induced harmful effects. The result is inconsistent with a number of previous studies which support the concept that NO can harm lung tissue in silica-induced effects by combining with superoxide to form peroxynitrite. We conclude that silica-induced harmful effects are probably caused by NO or peroxynitrite independent pathways in the rat model established by intratracheal instillation.

Methods

Experimental animals and group

The animals used in the experiments were male Wistar rats. They were purchased from the animal breeding facility of The University of New South Wales (Little Bay, NSW) and held in presterilised cages in the animal holding facility of the National Institute of Occupational Health and Safety. Sterilised food and water were supplied *ad libitum*. Experimental protocols involving rats were in accordance with the University of Sydney regulations and approved by The Animal Care Ethics Committee (ACEC) of The University of Sydney (ACEC number: WOR/7-95/2/2182).

In this study, body weights of the rats at the start of the experiment were 224–235 grams. The rats were grouped by randomised block design into four groups (each of 5): (1) it saline + ip saline; (2) it saline + ip L-NAME; (3) it silica + ip saline; (4) it silica + ip L-NAME.

Preparation of dust suspensions

Min-U-Sil 5 silica was obtained from Silica Corp, Berkeley Springs, West Virginia (particle size range was 0.6–8.0 µm, 98% < 5 µm diameter; purity was 99.5% α-quartz by X-ray diffraction). A 12.5 mg amount of dust was suspended in 0.5 ml saline or 0.5 ml saline alone, sterilised by autoclave and vortexed before injection. The preparation was free of endotoxin.

Intratracheal instillation and intraperitoneal injection

Rats were anaesthetised by ip injection of a mixture of ketamine (100 mg/kg) and xylazine (3.3 mg/kg, Sigma Chemical Company, St Louis, MO, USA). Before use, the chemicals were dissolved in saline and sterilised by filtration. Intratracheal instillation was performed by a procedure of tracheal exposure. After shaving, the skin in the ventral aspect of the neck was incised in the midline. The

trachea was exposed through blunt dissection. Using a 1 ml disposable syringe with a #26 needle, 12.5 mg silica dust dissolved in 0.5 ml of saline, or saline alone, was injected. The incision was sutured with interrupted silk sutures immediately after the injection and the rat was kept in a 30°C incubator until it regained consciousness. Immediately after the recovery of the instilled rats from anaesthesia, saline or L-NAME dissolved in saline was ip injected at a dosage of 15 mg/kg. Thereafter, the injection was conducted daily at the same dosage for 4 weeks.

Bronchoalveolar lavage

Randomly-selected rats were anaesthetised with an ip injection of 75 mg/kg of pentobarbital when 4 weeks observation period completed. Laparotomy was performed and the abdominal aorta was exposed. The rats were killed by transection of the aorta. There was no significant difference in body weight among the different groups of rats in each experiment when sacrificed.

After sacrifice, an incision was made into the neck region and the trachea was exposed by trimming of surrounding tissue. A piece of suturing thread was passed under the exposed trachea. A small opening was made by cutting in a lengthwise direction just below the larynx and in transverse direction just below and connected to the lengthwise incision. An 18-gauge needle in plastic tubing was placed into the opening and pushed through the trachea towards the lung. The needle was secured in place with the thread by ligation. A 5 ml volume of phosphate buffered saline (PBS) was injected into the needle with a 5 ml syringe for lavage. The injection process was finished within one minute and the fluid was kept in the lung for an additional minute. The withdrawal process was also finished in one minute. In total, two aliquots of 5 ml PBS were used for the lavage of each rat. The recovered fluid of each rat was pooled and the volume recorded. The recovered volume of BALF was between 9.0 to 9.7 ml and no significant difference could be detected among the different groups of rats in the recovered volume.

Counting of BALF cells

The total cell number was counted with a haemocytometer. A 50 µl volume of obtained lavage fluid was transferred to a plastic vial and mixed with the same amount of trypan blue solution (Sigma Chemical Company, St Louis, MO, USA). The well-mixed fluid then was transferred to a haemocytometer and the BALF cells were counted under a microscope.

Slide preparation, staining and differential counting

A 100 µl volume of the BALF was placed on a slide by Cytospin centrifugation immediately after the lavage (600 rpm, 5 min, Shandon, USA). At least two slides were prepared from each rat and stained with Diff-Quik (Lab. Aids

Pty. Ltd., Narrabeen, Australia). The interval between sacrifice of rat to placement of BALF on slides was approximately 15 min.

The slides were read under oil immersion ($\times 1000$). Five hundred leucocytes were counted to determine the frequency of different types of cells by their morphology.

Measurement of BALF protein

The supernatant of centrifuged BALF was placed in 400 µl polyethylene tubes, in duplicate. The protein concentration was measured by Lowry's method [44] using bovine serum albumin as a standard. The measurement was conducted 1 day after lavage.

Measurement of lactate dehydrogenase

BALF was centrifuged (3000 rpm, 15 min) and the supernatant was used for measurement of LDH activity. The measurement was performed with a CentrifChem System 400 using a kit method (Trace Scientific Pty. Ltd., Sydney, Australia) and determined within 3 days after lavage. LDH measurements were performed within 3 days after lavage. The samples were kept in ice during operation and stored in -20°C freezer.

Identification of apoptosis

The slides were blind-coded before the scoring of apoptotic leucocytes. A minimum of 1000 leucocytes were counted for the occurrence of cells with apoptotic features. A minimum of 500 neutrophils were counted to determine the percentage of apoptotic neutrophils. Apoptotic features included formation of condensed chromatin bodies with sharp edges and convolution of the cell surface [45,46]. Apoptotic macrophages and neutrophils were differentiated by morphology. Apoptotic macrophages were generally larger in size, with clear and non-ciliated cell borders. The key point in discriminating apoptotic macrophages from apoptotic neutrophils is the cytoplasm staining with Diff-Quik. The cytoplasm of apoptotic macrophages is a blue colour, similar to the non-apoptotic macrophages. The cytoplasm of apoptotic neutrophils is light yellow, similar to non-apoptotic neutrophils. Morphological apoptosis of the cells in BALF was confirmed by agarose gel electrophoresis of extracted genomic DNA. In the electrophoresis, cells in BALF with morphological apoptosis showed a ladder pattern, whereas cells without morphological apoptosis lacked such an electrophoretic feature.

Scoring of macrophages with engulfed apoptotic cells

A total of 1000 macrophages were counted for the incidence of macrophages with engulfed cells or apoptotic bodies. The proportion of macrophages with engulfed cells or bodies in 1000 counted macrophages was calculated.

Effect of L-NAME injection on production of NO metabolites

To examine the effects of L-NAME administration on production of NO in silica-instilled lungs, forty male Wistar rats were used to obtain BALF for the measurement of nitrite/nitrate levels. Body weights of these rats at the start of the experiment were 209–220 grams. The rats were grouped by randomised block design into four groups for intratracheal instillation and ip injection (each of 10): (1) it saline + ip saline; (2) it saline + ip L-NAME; (3) it silica + ip saline; (4) it silica + ip L-NAME. At two weeks after instillation, 5 rats in each group were randomly selected and killed for lavage. At four weeks after instillation, all the remaining rats were killed and lavaged. The lavage fluid was used for the measurement of nitrite/nitrate level within two hours. The procedures in instillation, injection, killing and lavage were exactly the same as described above.

Nitrite/nitrate levels in the obtained BALF were measured by a Total Nitric Oxide and Nitrate/Nitrite Parameter Assay Kit (R&D System, Minneapolis MN, USA) with conversion of nitrate to nitrite by nitrate reductase. Absorbances were measured at 540 nm with ELISA microplate reader (Titertek Multiskan MCC/340).

Statistical analysis

The results are expressed as means \pm standard error (se). Student-Newman-Keul's test was used for multiple comparison of different animal groups of each index with the software package InStat because this test method is suitable for multiple comparison. Analyses of variance were also conducted by this software. Statistical significance was preset at $p < 0.05$.

Competing Interests

The author(s) declare that they have no competing interests.

Authors' contributions

HW designed the study, conducted all the experiments and drafted the manuscript.

JL participated in the design of the study, contributed to the discussion and provided funding for the project.

All authors read and approved the final manuscript.

Acknowledgements

The authors would like to thank Dr Tony Bonin for useful discussion during the implementation of this study. This study was partially supported by Dust Diseases Board Research Grants Scheme.

References

- Fubini B, Hubbard A: **Reactive oxygen species (ROS) and reactive nitrogen species (RNS) generation by silica in inflammation and fibrosis.** *Free Radic Biol Med* 2003, **34**:1507-1516.
- Zeidler P, Hubbs A, Battelli L, Castranova V: **Role of inducible nitric oxide synthase-derived nitric oxide in silica-induced pulmonary inflammation and fibrosis.** *J Toxicol Environ Health A* 2004, **67**:1001-1026.
- Blackford JA Jr., Antonini JM, Castranova V, Dey RD: **Intratracheal instillation of silica up-regulates inducible nitric oxide synthase gene expression and increases nitric oxide production in alveolar macrophages and neutrophils.** *Am J Respir Cell Mol Biol* 1994, **11**:426-431.
- Castranova V, Huffman LJ, Judy DJ, Bylander JE, Lapp LN, Weber SL, Blackford JA, Dey RD: **Enhancement of nitric oxide production by pulmonary cells following silica exposure.** *Environ Health Perspect* 1998, **106 Suppl 5**:1165-1169.
- Davis GS, Pfeiffer LM, Hemenway DR: **Interferon-gamma production by specific lung lymphocyte phenotypes in silicosis in mice.** *Am J Respir Cell Mol Biol* 2000, **22**:491-501.
- Feng HM, Walker DH: **Interferon-gamma and tumor necrosis factor-alpha exert their antirickettsial effect via induction of synthesis of nitric oxide.** *Am J Pathol* 1993, **143**:1016-1023.
- Naseem KM, Bruckdorfer KR: **Hydrogen peroxide at low concentrations strongly enhances the inhibitory effect of nitric oxide on platelets.** *Biochem J* 1995, **310 (Pt 1)**:149-153.
- Tada Y, Mori T, Shinogi T, Yao N, Takahashi S, Betsuyaku S, Sakamoto M, Park P, Nakayashiki H, Tosa Y, Mayama S: **Nitric oxide and reactive oxygen species do not elicit hypersensitive cell death but induce apoptosis in the adjacent cells during the defense response of oat.** *Mol Plant Microbe Interact* 2004, **17**:245-253.
- Leigh J, Wang H, Bonin A, Peters M, Ruan X: **Silica-induced apoptosis in alveolar and granulomatous cells in vivo.** *Environ Health Perspect* 1997, **105 Suppl 5**:1241-1245.
- Terenzi F, Diaz-Guerra MJ, Casado M, Hortelano S, Leoni S, Bosca L: **Bacterial lipopeptides induce nitric oxide synthase and promote apoptosis through nitric oxide-independent pathways in rat macrophages.** *J Biol Chem* 1995, **270**:6017-6021.
- Washo-Stultz D, Hoglen N, Bernstein H, Bernstein C, Payne CM: **Role of nitric oxide and peroxynitrite in bile salt-induced apoptosis: relevance to colon carcinogenesis.** *Nutr Cancer* 1999, **35**:180-188.
- Beckman JS, Beckman TW, Chen J, Marshall PA, Freeman BA: **Apparent hydroxyl radical production by peroxynitrite: implications for endothelial injury from nitric oxide and superoxide.** *Proc Natl Acad Sci U S A* 1990, **87**:1620-1624.
- Koppenol WH, Moreno JJ, Pryor WA, Ischiropoulos H, Beckman JS: **Peroxynitrite, a cloaked oxidant formed by nitric oxide and superoxide.** *Chem Res Toxicol* 1992, **5**:834-842.
- Thomas G, Ando T, Verma K, Kagan E: **Asbestos fibers and interferon-gamma up-regulate nitric oxide production in rat alveolar macrophages.** *Am J Respir Cell Mol Biol* 1994, **11**:707-715.
- Pendino KJ, Laskin JD, Shuler RL, Punjabi CJ, Laskin DL: **Enhanced production of nitric oxide by rat alveolar macrophages after inhalation of a pulmonary irritant is associated with increased expression of nitric oxide synthase.** *J Immunol* 1993, **151**:7196-7205.
- Walley KR, McDonald TE, Higashimoto Y, Hayashi S: **Modulation of proinflammatory cytokines by nitric oxide in murine acute lung injury.** *Am J Respir Crit Care Med* 1999, **160**:698-704.
- Hogaboam CM, Chensue SW, Steinhauser ML, Huffnagle GB, Lukacs NW, Strieter RM, Kunkel SL: **Alteration of the cytokine phenotype in an experimental lung granuloma model by inhibiting nitric oxide.** *J Immunol* 1997, **159**:5585-5593.
- Hogaboam CM, Gallinat CS, Bone-Larson C, Chensue SW, Lukacs NW, Strieter RM, Kunkel SL: **Collagen deposition in a non-fibrotic lung granuloma model after nitric oxide inhibition.** *Am J Pathol* 1998, **153**:1861-1872.
- Nunes H, Lebrech D, Mazmanian M, Capron F, Heller J, Tazi KA, Zerbib E, Dulmet E, Moreau R, Dinh-Xuan AT, Simonneau G, Herve P: **Role of nitric oxide in hepatopulmonary syndrome in cirrhotic rats.** *Am J Respir Crit Care Med* 2001, **164**:879-885.
- Antonini JM, Van Dyke K, Ye Z, DiMatteo M, Reasor MJ: **Introduction of luminol-dependent chemiluminescence as a method to study silica inflammation in the tissue and phagocytic cells of rat lung.** *Environ Health Perspect* 1994, **102 Suppl 10**:37-42.

21. Maeda H, Akaike T: **Nitric oxide and oxygen radicals in infection, inflammation, and cancer.** *Biochemistry (Mosc)* 1998, **63**:854-865.
22. Lee J: **Inducible nitric oxide synthase mRNA expression and nitric oxide production in silica-induced acute inflammatory lung injury.** *Korean-Journal-of-Physiology-and-Pharmacology* 1998, **2**:233-239.
23. Brune B, Mohr S, Messmer UK: **Protein thiol modification and apoptotic cell death as cGMP-independent nitric oxide (NO) signaling pathways.** *Rev Physiol Biochem Pharmacol* 1996, **127**:1-30.
24. Hu J, Van Eldik LJ: **S100 beta induces apoptotic cell death in cultured astrocytes via a nitric oxide-dependent pathway.** *Biochim Biophys Acta* 1996, **1313**:239-245.
25. Aliprantis AO, Diez-Roux G, Mulder LC, Zychlinsky A, Lang RA: **Do macrophages kill through apoptosis?** *Immunol Today* 1996, **17**:573-576.
26. Srivastava KD, Rom WN, Jagirdar J, Yie TA, Gordon T, Tchou-Wong KM: **Crucial role of interleukin-1beta and nitric oxide synthase in silica-induced inflammation and apoptosis in mice.** *Am J Respir Crit Care Med* 2002, **165**:527-533.
27. Wiethoff AJ, Reed KL, Webb TR, Warheit DB: **Assessing the role of neutrophil apoptosis in the resolution of particle-induced pulmonary inflammation.** *Inhal Toxicol* 2003, **15**:1231-1246.
28. Bosca L, Hortelano S: **Mechanisms of nitric oxide-dependent apoptosis: involvement of mitochondrial mediators.** *Cell Signal* 1999, **11**:239-244.
29. Rogers MJ, Chilton KM, Coxon FP, Lawry J, Smith MO, Suri S, Russell RG: **Bisphosphonates induce apoptosis in mouse macrophage-like cells in vitro by a nitric oxide-independent mechanism.** *J Bone Miner Res* 1996, **11**:1482-1491.
30. Mitrovic B, Ignarro LJ, Vinters HV, Akers MA, Schmid I, Uittenbogaart C, Merrill JE: **Nitric oxide induces necrotic but not apoptotic cell death in oligodendrocytes.** *Neuroscience* 1995, **65**:531-539.
31. Delaney CA, Tyrberg B, Bouwens L, Vaghef H, Hellman B, Eizirik DL: **Sensitivity of human pancreatic islets to peroxynitrite-induced cell dysfunction and death.** *FEBS Lett* 1996, **394**:300-306.
32. Estevez AG, Radi R, Barbeito L, Shin JT, Thompson JA, Beckman JS: **Peroxynitrite-induced cytotoxicity in PC12 cells: evidence for an apoptotic mechanism differentially modulated by neurotrophic factors.** *J Neurochem* 1995, **65**:1543-1550.
33. Ye Y, Estevez AG: **Nitrotyrosine containing proteins in PC12 cells after exposure to peroxynitrite.** *Society for Neuroscience Abstract Viewer and Itinerary Planner* 2002, **303**:17.
34. Shin JT, Barbeito L, MacMillan-Crow LA, Beckman JS, Thompson JA: **Acidic fibroblast growth factor enhances peroxynitrite-induced apoptosis in primary murine fibroblasts.** *Arch Biochem Biophys* 1996, **335**:32-41.
35. Zhuang S, Simon G: **Peroxynitrite-induced apoptosis involves activation of multiple caspases in HL-60 cells.** *Am J Physiol Cell Physiol* 2000, **279**:C341-51.
36. Lin KT, Xue JY, Nomen M, Spur B, Wong PY: **Peroxynitrite-induced apoptosis in HL-60 cells.** *J Biol Chem* 1995, **270**:16487-16490.
37. Cox G: **IL-10 enhances resolution of pulmonary inflammation in vivo by promoting apoptosis of neutrophils.** *Am J Physiol* 1996, **271**:L566-71.
38. Shimizu S, Nomoto M, Yamamoto T, Momose K: **Reduction by NG-nitro-L-arginine of H2O2-induced endothelial cell injury.** *Br J Pharmacol* 1994, **113**:564-568.
39. Seif-el-Nasr M, Fahim AT: **Antioxidant effect of N omega-nitro-L-arginine methyl ester (L-NAME) on global cerebral ischemia in a rat model.** *Arzneimittelforschung* 2001, **51**:628-632.
40. Wishah K, Malur A, Raychaudhuri B, Melton AL, Kavuru MS, Thomasen MJ: **Nitric oxide blocks inflammatory cytokine secretion triggered by CD23 in monocytes from allergic, asthmatic patients and healthy controls.** *Ann Allergy Asthma Immunol* 2002, **89**:78-82.
41. Peng HB, Spiecker M, Liao JK: **Inducible nitric oxide: an autoregulatory feedback inhibitor of vascular inflammation.** *J Immunol* 1998, **161**:1970-1976.
42. Dorger M, Allmeling AM, Kiefmann R, Schropp A, Krombach F: **Dual role of inducible nitric oxide synthase in acute asbestos-induced lung injury.** *Free Radic Biol Med* 2002, **33**:491-501.
43. Zeidler PC, Castranova V: **Role of nitric oxide in pathological responses of the lung to exposure to environmental/occupational agents.** *Redox Rep* 2004, **9**:7-18.
44. Lowry OH, Rosebrough NJ, Farr AL, Randall RJ: **Protein measurement with the Folin phenol reagent.** *J Biol Chem* 1951, **193**:265-275.
45. Newman SL, Henson JE, Henson PM: **Phagocytosis of senescent neutrophils by human monocyte-derived macrophages and rabbit inflammatory macrophages.** *J Exp Med* 1982, **156**:430-442.
46. Savill JS, Wyllie AH, Henson JE, Walport MJ, Henson PM, Haslett C: **Macrophage phagocytosis of aging neutrophils in inflammation. Programmed cell death in the neutrophil leads to its recognition by macrophages.** *J Clin Invest* 1989, **83**:865-875.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

