

	<u>Dominula</u>	<u>Heterozygotes</u>	<u>Bimacula</u>	<u>Total</u>	<u>Gene frequency %</u>
up to 1928	164	4 (2.4%)	-	168	1.2
1939	184	37 (16.6%)	2	223	9.8 1.2
1940	92	24 (20.5%)	1	117	11.1
1941	400	59 (12.8%)	2	461	6.8
1942	183	22 (10.7%)	-	205	5.4
1943	239	30 (11.2%)	-	269	5.6
1944	452	43 (8.7%)	1	496	4.5
1945	326	44 (11.8%)	2	372	6.5

Data received from Dr. Ford, March 1946.

	<u>dominula</u>	<u>medionigra</u>	<u>bimaculata</u>	<u>Total.</u>	<u>Sex-frequency</u> <u>%</u>
1928	164	4 (2.4%)	-	168	1.2
1939	184	37 (16.6%)	2	223	9.8
1940	92	24 (20.5%)	1	117	11.1
1941	400	59 (12.8%)	2	461	6.8
1942	183	22 (10.7%)	-	205	5.4

1943.

	<u>dominula</u>	<u>mediogra</u>	<u>bi macula</u>	<u>Total.</u>	
9. VII.	15	3	-	18	
10. VII.	13	1	-	14	(including 1 dead in spiders web not included in Δ).
11. VII.	20	2	-	22	
12. VII.	21	3	-	24	
14. VII.	34	5	-	39	
15. VII.	38	2	-	40	
18. VII.	38	3	-	41	
19. VII.	17	3	-	20	
					$\frac{22}{218}$

