

PUBLISHED VERSION

Nick Schofield, Craig Magee, Simon Holford, and Christopher Jackson
Magma plumbing and emplacement mechanisms within sedimentary basins
Geophysical Research Abstracts, Vol. 15, 2013; EGU2013-2251

© Author(s) 2013. CC Attribution 3.0 License.

PERMISSIONS

<http://creativecommons.org/licenses/by/3.0/>



This is a human-readable summary of (and not a substitute for) the [license](#).

[Disclaimer](#)



You are free to:

Share — copy and redistribute the material in any medium or format

Adapt — remix, transform, and build upon the material

for any purpose, even commercially.

The licensor cannot revoke these freedoms as long as you follow the license terms.

Under the following terms:



Attribution — You must give **appropriate credit**, provide a link to the license, and **indicate if changes were made**. You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.

No additional restrictions — You may not apply legal terms or **technological measures** that legally restrict others from doing anything the license permits.

<http://hdl.handle.net/2440/84342>



Magma Plumbing and Emplacement Mechanisms within Sedimentary Basins

Nick Schofield (1), Craig Magee (2), Simon Holford (3), and Christopher Jackson (2)

(1) Earth Sciences, School of Geography, Earth and Environmental Sciences, University of Birmingham, Edgbaston, Birmingham, B15 2TT, England (n.schofield@bham.ac.uk), (2) Department of Earth Science and Engineering, Imperial College, Prince Consort Road, London, SW7 2BP, England, UK, (3) Australian School of Petroleum Centre for Tectonics, Resources and Exploration (TRaX) University of Adelaide, Adelaide SA 5005

In recent years our understanding of sub-volcanic magmatic plumbing systems has been revolutionised by the study of hydrocarbon industry 3D seismic reflection datasets from offshore sedimentary basins. In particular, 3D seismic reflection data has provided important insights into sheet intrusion geometry and emplacement mechanisms as well as linkages and magma flow between multiple intrusions within sill-complexes. However, even high-quality 3D seismic reflection datasets have a limit to what they can resolve; thus, to allow a better understanding of detailed emplacement mechanisms and to test the validity of subsurface-based interpretations, it is critical to bridge the resolution gap that exists between seismic and outcrop datasets.

Magmatic sheet (sill) intrusions contribute significantly to the upper crustal magma transport network. The emplacement mechanism of the magmatic sheets controls the final geometry of the intrusions and the characteristics of host rock deformation. Previous observations have highlighted the preponderance of brittle structures (e.g. intrusive steps and broken brides) associated with shallow-level sheet intrusions. However, recent studies have suggested that non-brittle host rock behaviour also occurs, particularly related to the formation of magma fingers during shallow-level sill intrusion. Importantly, these structures can provide insights into emplacement style and magma flow directions. Here, we examine both brittle and non-brittle intrusion mechanisms and structures using both field- and 3D seismic-based observations from a series of widespread and variable magmatic systems. Non-brittle emplacement (i.e. magma finger and lobe development) appears to be primarily associated with viscous flow of the host rock during intrusion and is therefore intimately linked to the contemporaneous host rock rheology as well as magma dynamics. Purely brittle and non-brittle emplacement processes are found to be end members with many intrusions containing evidence of both behaviours.

Deriving the host rock characteristics is therefore an important step in discerning potential diagnostic magma flow indicators and intrusion geometries within field-, seismic- and model-based studies. Incorporation of variable host material behaviours in numerical and analogue modelling, tuned using direct field observations, may consequently further our understanding of the controls on shallow-level intrusion.