Identification of Sea Breezes, their Climatic Trends and Causation, with Application to the Adelaide Coast

ZAHRA PAZANDEH MASOULEH

BENG (CIVIL & STRUCTURAL), MSCENG (Coastal)

A THESIS SUBMITTED FOR THE DEGREE OF DOCTOR OF PHILOSOPHY



SCHOOL OF CIVIL, ENVIRONMENTAL AND MINING ENGINEERING July 2015

Table of contents

Abstract		
Statement of Originality	iv	
Acknowledgement	v	
1. Introduction	1	
1.1. Background and motivation for the research	1	
1.2. Research scope and Objective	2	
1.3. Overview of Thesis	5	
2. Literature Review	7	
2.1. Introduction to Adelaide Coastal Climate	7	
2.2. Sea breeze Circulation System	8	
2.3. Geophysical variable that affect sea breezes	9	
2.4. Surface energy budget	12	
2.5. Urban Boundary Layers	13	
2.6. Thermal Properties and the Urban Street Canyon Effect	15	
2.7. Urban Heat Island	16	
2.7.1. Heat Island Types and Circulation	18	
2.7.2. Urban heat island impact	19	
2.7.3. Characteristics of Urban Heat Island in Different Cities Worldwide	20	
2.7.4. Interaction of Urban Heat Island and Sea breeze	22	
2.8. Sea Breeze Detection Methods	24	
2.9. Numerical Simulation of climate	26	
2.9.1. Regional Climate Models (RCMs)		
2.9.2. Application of RCMs		
2.9.2.1. Simulation of Sea-land breeze		
2.9.2.2. Simulation of Land cover change and urban Climate		
2.9.3. Weather Research and Forecasting Model (WRF)		

2.9.3.1. The major feature of ARW and the governing equations	
3. Climate of the Study Area	41
3.1. Description of the Adelaide Metropolitan Area	41
3.2. The main feature of climate in Adelaide	41
3.2.1. Wind climate	
3.3. South Australian Major Climate Drivers	44
3.3.1. The El Niño Southern Oscillation	44
3.3.2. The Indian Ocean Dipole	45
3.3.3. Southern Annular Mode	45
3.3.4. Sub-Tropical Ridge	46
3.3.5. Cloud bands	46
3.3.6. Frontal Systems	47
3.3.7. Cut-Off Lows	47
3.3.8. Blocking Highs	47
3.4. Previous study of sea breezes in the region	
4. Adelaide Sea breeze	50
4.1. Adelaide Mesoscale Meteorology	50
4.2. Detection and Validation of Sea Breezes	51
4.2.1. Adelaide Airport	
4.2.2. Edithburgh	54
4.2.3. The Sea Breeze Identification algorithm	55
4.2.4. Method Assumptions	57
4.3. The Results	57
4.4. Sensitivity analysis	59
4.4.1. Sea surface temperature	59
4.4.2. Sensitivity to land surface temperature	60
4.4.3. Sensitivity to surface level wind speed criteria	61
	<i></i>

4.6. Long-term trend	64
4.6.1. Some considerations on statistical analysis of data	64
4.6.2. Summer	
4.6.3. Autumn	71
4.6.4. Spring	74
4.7. Daily cycle of wind	77
4.8. Analysis and discussion	
4.8.1. Effect of Change to the Land surface temperature	
4.8.2. Effect of Other Climate Drivers	
4.8.3. Effect of local climate drivers and urban heat island	
5. Numerical Modelling of the urban climate	
5.1. WRF model principles	
5.2. Models initial and boundary conditions	
5.3. Simulation set-up	95
5.4. Validation of model	
5.4.1. Statistical Metrics	
5.5. Modelling Metropolitan with pre-European Land cover	
5.5.1. Temperature	
5.5.2. Wind Speed	
5.6. Discussion	
6. Summary and conclusion	
7. References	
Appendix A: Adelaide Airport Station Metadata	
Appendix B: Climatic Indices and their Correlation with monthly a components of wind	averaged 192

List of Figures

FIGURE 1.1. THE LOCATION OF METROPOLITAN ADELAIDE (GOOGLE EARTH)
FIGURE 2.1. GEOPHYSICAL VARIABLE THAT CONTROL SEA AND LAKE BREEZES (CORIOLIS PARAMETER F NOT
SHOWN) ADAPTED FROM CROSMAN AND HOREL (2010)
FIGURE 2.2. THE URBAN LAYER STRATIFICATIONS (AFTER OKE, 1988)
FIGURE 2.3. THE THERMAL DIFFERENTIAL OF CITY AND ITS ADJACENT AREAS (GOODMAN, 1999)
FIGURE 2.4. A SCHEMATIC DIAGRAM OF URBAN PLUM AND THE HEAT ISLAND CIRCULATION
FIGURE 2.5. TEMPERATURE DISTRIBUTION IN DAYTIME OF 13 OF AUGUST 2009, OKAYAMA CITY IN JAPAN -
DARK GREY AND BROWN SHADING REPRESENTS COMMERCIAL AREAS AND MOUNTAIN RESPECTIVELY
(SHIGETA <i>ET AL</i> , 2009)
FIGURE 2.6. AN EXAMPLE OF THE LONGEST TRAJECTORIES THAT ORIGINATED UPWIND (OPEN CIRCLES) AND
DOWNWIND (FULL CIRCLES) OF THE PLUME AXIS IN THE PRESENCE OF A SEA BREEZE, TAKEN FROM A
STUDY BY CENEDESE AND MONTI (2003), $x/D = 0$ represents the centre of the urban area 23
FIGURE 2.7. AN EXAMPLE OF NESTING IN A REGIONAL CLIMATE MODEL WITH THE HORIZONTAL GRID SHOWN
IN BLACK LINE, BOUNDARY OF BUFFER ZONE IN RED LINE AND VERTICAL COLUMN INDICATING THE
ATMOSPHERIC LAYERS BY THE MODEL, FROM RASCH (2012)
FIGURE 2.8. THE INTERACTIONS BETWEEN THE FIVE MAJOR PHYSICAL PARAMETERIZATIONS USED IN THE. 35
WRF-ARW (Dudhia, 2010)
FIGURE 3.1.A. SEASONAL WIND ROSE IN ADELAIDE FOR SUMMER, 9 AM ON LEFT AND 3 PM ON RIGHT
(AUSTRALIAN BUREA OF METEOROLOGY, 2014A)
FIGURE 3.1.B. SEASONAL WIND ROSE IN ADELAIDE, 9 AM ON LEFT AND 3 PM ON RIGHT (AUSTRALIAN BUREA
OF METEOROLOGY, 2014A) 43
FIGURE 3.2. CLIMATIC DRIVERS OF AUSTRALIA (AUSTRALIAN BUREAU OF METEOROLOGY, 2010)
FIGURE 4.1. MEAN RATIOS CGA/PTA FOR MEAN WIND SPEED (\overline{R}) and GUST WIND (\overline{R}) of day (solid
LINE) AND NIGHT (PECKED LINE) AND ALL OBSERVATION (DOTTED LINE), FROM (LOGUE, 1986) 53
FIGURE 4.2. LOCATION OF ADELAIDE AIRPORT AND EDITHBURGH STATIONS
FIGURE 4. 3. THE FREQUENCY OF SEA BREEZE EVENT FOR EACH SEASON, THE BOX SHOWS THE LOWER AND
THE UPPER QUARTILE
FIGURE 4.4. AVAILABLE (GREEN DOTS) AND AVERAGED (RED LINE) SEA SURFACE TEMPERATURE AGAINST
THE OBSERVED LAND SURFACE TEMPERATURE FOR EACH DAY OF THE YEAR
FIGURE 4.5. WIND HODOGRAPH OF SEA BREEZE (FULL LINE) AND NON-SEA BREEZE DAYS (DOTTED LINE) FOR
BOTH SIDES OF THE GULF (1993-2008). THE CIRCLES SHOW THE WIND SPEED AT THE INTERVAL OF 1
M/S
FIGURE 4.6. AVERAGE SEA BREEZE DAYS' U-COMPONENT (A) AND V-COMPONENT (C) OF ADELAIDE
AIRPORT (SOLID LINE) AND EDITHBURGH (DASHED LINE). SAME FOR NON-SEA BREEZE DAYS (B) AND
(D) RESPECTIVELY
FIGURE 4.7. U COMPONENT OF 15:00 HOUR FOR ENTIRE PERIOD (A) AND EXCLUDING 1972-1984 (B) 66
FIGURE 4.8. U COMPONENT OF 12:00 HOUR FOR ENTIRE PERIOD (A) AND EXCLUDING 1972-1984 (B) AND
SIMILARLY FOR 21:00 DATA (C,D)
Figure 4.9. V component of $18:00~({\rm a})$ and $21:00~({\rm b})$ winds of sea breeze days and $21:00~({\rm c})$ wind of
NON-SEA BREEZE DAYS
Figure 4.10. U component of 15:00 (a) and 21:00 (b) wind of sea breeze and 21:00 (c) wind of non-
SEA BREEZE DAYS. THE GREY LINE SHOWS THE FIVE YEARS MOVING AVERAGE
FIGURE 4.11. V COMPONENT OF SEA BREEZE (LEFT) AND NON-SEA BREEZE (RIGHT) AT 15:00 (A,B), 18:00
(C,D) AND 21:00 (E,F)
FIGURE 4.12. THE U COMPONENT OF SEA BREEZE DAY 12:00 (A), 15:00 (B), 18:00 (C) AND 21:00 (D) WIND.
FIGURE 4.13. THE V COMPONENT OF SEA BREEZE DAY 15:00 (A), 18:00 (B) AND 21:00 (C) WIND

FIGURE 4.14. SEA BREEZE (SOLID LINE) AND NON-SEA BREEZE DAYS (DASHED LINE) HODOGRAPH (1955-2007) FOR ADELAIDE AIRPORT. THE SEA BREEZE SECTOR (180°-320°) IS ILLUSTRATED AS A BLUE
CURVE. THE CIRCLES SHOW THE WIND SPEED AT THE INTERVAL OF 1 M/S
FIGURE 4.15. THE FREQUENCY OF TIME OF SEA BREEZE START (A) CESSATION (B) AND MAXIMUM (C) (1985-2007) IN EACH SEASON
FIGURE 4.16. MONTHLY AVERAGES OF SEA BREEZE START-END AND MAXIMUM (1985-2007)
FIGURE 4.17. AVERAGE OF MAXIMUM SEA BREEZE DIRECTION AND SPEED (1985-2007). THE NUMBERS ON THE ARROWS ARE THE WIND SPEED IN M S^{-1}
FIGURE 4.18. PERCENTAGE OF ANNUAL SEA BREEZE EVENTS FOR THE PERIOD OF 1956-2007, THE LINE
SHOWS THE 5-YEAR MOVING AVERAGE
FIGURE 4.19. THE 18:00 (X AXES) AGAINST 15:00 (Y AXES) WIND COMPONENT FOR SEA BREEZE DAYS U(A) $V_{c}(c)$ component (M s ⁻¹) and shall and y for non-sea prefere days (n, n)
, $V(C)$ COMPONENT (M.S.) AND SIMILARL 1 FOR NON-SEA BREEZE DATS (B, D)
FIGURE 4.20. ANOMALY OF 1.2 M MAXIMUM (A) AND MINIMUM (B) TEMPERATURES IN ADELAIDE AIRPORT. 84
FIGURE 4.21. PLOT OF MAXIMUM TEMPERATURE AGAINST MONTHLY AVERAGED V COMPONENTS OF WIND
ON SEA BREEZE DAYS, AT $18:00$ (A) and $21:00(B)$ and non-sea breeze days (C, d respectively).
FIGURE 4.22. PLOT OF MONTHLY AVERAGED MAXIMUM TEMPERATURE AND THE PERCENTAGE OF SELECTED
SEA BREEZE DAYS
FIGURE 4.23. AVERAGED SOUTHERLY WIND COMPONENT OF JANUARY 18:00 (A) AND 21:00 (B) WITH
ACTUAL VALUE (GRAY DOTS) AND DE-TRENDED VALUE (BLACK DOTS)
FIGURE 4.24. ADELAIDE METROPOLITAN POPULATION (AUSTRALIAN BUREAU OF STATISTICS, 2012)
FIGURE 4.25. ADELAIDE CITY EXPANSION SINCE 1910 (STATE OF ENVIRONMENT, 2011) ON RIGHT AND
POPULATION DENSITY IN 2012 (ABS, 2013) ON LEFT
FIGURE 5.1. LAYOUT OF MODEL DOMAINS, D1, D2 AND D3, WITH ENLARGED DOMAIN 3 SHOWN ON THE
RIGHT (GOOGLE EARTH, 2013)
FIGURE 5.2. SCHEMATIC OF SINGLE-LAYER URBAN CANOPY MODEL (FROM CHEN ET AL., 2011)
FIGURE 5.5. THE CHANGE OF VEGETATION IN SOUTH AUSTRALIA SINCE 1/50 (DEPARTMENT OF THE
ENVIRONMENT AND WATER RESOURCES, 2007)
FIGURE 5.4. LAND COVER INDEX FROM THE MODEL FOR CTL SIMULATION (LEFT) AND INT V (RIGHT)
PERIOD OF SIMULATION (DISPLAYED BY RED FLAGS) OF AVAILABLE METEOROLOGICAL STATIONS FOR THE PERIOD OF SIMULATION (GOOGLE EARTH, 2013), ADELAIDE METROPOLITAN AREA IS SHOWN BY
TELLOW LINE
TIGURE J.U. TATLOR DIAGRAM OF STATISTICAL VALUES FOR ALL STATIONS FOR THREE VARIABLES. TEMDED ATUDE (CIDCLES) WIND SPEED (SOLIADES) AND WIND DIRECTION (TRIANCLES) 106
FIGURE 5.7. THE PERFORMANCE OF MODEL TO SIMULATE 2M TEMPERATURE (A), 10M WIND SPEED (B) AND
10M WIND DIRECTION (C) IN ADELAIDE AIRPORT IN FEBRUARY 2005. BLUE DOTS REPRESENT THE
OBSERVATION AND BLACK LINES ARE THE MODEL'S OUTPUTS
FIGURE 5.8. THE AVERAGED DAILY CYCLE OF TEMPERATURE SIMULATED WITH CURRENT LAND COVER
(BLACK) AND NATIVE VEGETATION (BLUE) AT AN INNER SUBURB (A), A COASTAL POINT (B) AND THE DIFFERENCE $(C) = (A)_{-}(B)$
FIGURE 5.9 THE DIFFERENCE RETWEEN AN HOURLY AVERAGED TEMPERATURE OF AN INNER CITY LOCATION
OF TWO SIMULATION ($T_{cTT} - T_{VTTV}$) FOR TWO COLD MONTHS OF MAY AND JULY AND TWO WARM
MONTHS OF JANUARY AND MARCH
FIGURE 5.10 Two examples of 2m height temperature ($^{\circ}C$) distribution over the area for two
TIMES STEPS: CTL (LEFT) AND NTV (RIGHT).
FIGURE 5.11. AREA AVERAGE WIND SPEED AT 10M HEIGHT OVER THE METROPOLITAN AREA. CTL (BLACK
DOTS AND THIN BLACK LINE) AND NTV (RED DOTS AND THIN GREY LINE) FOR THE PERIOD OF 13 TO 25
OF JANUARY 2005
FIGURE 5.12. SAME AS FIGURE 5.11 FOR AN ADJACENT POINT OVER WATER (GREEN AND BLUE OVALS
DEMONSTRATE THE DIFFERENCE BETWEEN TWO SIMULATIONS IN NIGHT TIMES AND AFTERNOON
RESPECTIVELY)
FIGURE 5.13. THE MONTHLY AVERAGED U-WIND COMPONENT OF THE WATER POINT AT LOWEST 2 KM LEVEL
OF WARM MONTHS (LINES) AND COLD MONTHS (DASHED LINE) AT TIMES OF 0600, 0900, 1200, 1500,

1800 and 2100 . The red and blue lines demonstrate the average of warm and cold
MONTHS, RESPECTIVELY
FIGURE 5.14. SAME AS FIGURE 5.13 FOR V-WIND COMPONENT
FIGURE 5.15. THE WARM PERIOD TIME-AVERAGED U (TWO LEFT COLUMNS) AND V (TWO RIGHT COLUMNS)
COMPONENT OF THE WATER POINT SIMULATED WITH CTL (black dots and thin line) and NTV
(BLUE DOTS AND THIN DASHED LINE) AT TIMES OF 0600 AND 0900 (TOP ROW), 1200AND 1500 (MIDDLE
ROW), 1800 AND 2100 (BOTTOM ROW)
FIGURE 5.16. THE HODOGRAPH OF TIME-AVERAGED WIND OF WMP AT 10M HEIGHT OF THE WATER POINT
FOR CTL (BLACK DOTS AND LINE) AND NTV (BLUE DOTS AND LINE) SIMULATIONS. THE CIRCLES
SHOW THE WIND SPEED INTERVALS OF 1 m s $^{-1}$ and the numbers denote the Australian Central
STANDARD TIME. AN EXAMPLE OF AVERAGED WIND VECTOR FOR THE HOUR OF 1400 in CTL run is
DEMONSTRATED BY AN ARROW
FIGURE 5.17. THE AVERAGED U-WIND COMPONENT PROFILE OF WMP IN 9 AM (ABOVE) AND 9 PM (BELOW)
FROM SHORELINE TO FURTHER 27 KM OVER WATER
FIGURE 5.18. THE VERTICAL COMPONENT OF WINDS SIMULATED WITH CTL (BLACK) AND NTV (BLUE) AT
TIMES OF 0000, 0300, 0600, 0900, 1200, 1500, 1800 and 2100 124
FIGURE 5.19. HODOGRAPH OF AVERAGED SEA BREEZE (LEFT) AND NON-SEA BREEZE DAY (RIGHT) OF WMP,
SIMULATED WITH CTL (BLACK) AND NTV (BLUE). THE CIRCLES SHOW THE WIND SPEED INTERVALS OF
1 m s ⁻¹ and the numbers denote the Australian Central Standard Time 125

List of Tables

TABLE 3.1: CLIMATIC DRIVERS OF SOUTH AUSTRALIA 48
TABLE 3.2: A SUMMARY OF SEA BREEZE DAY CHARACTERISTIC ADOPTED FROM PHYSICK AND BYRON-
Scott (1977)
TABLE 4.1: ADELAIDE AIRPORT STATION. DATA SUPPLIED BY AUSTRALIAN BUREA OF METEOROLOGY
(2014A)
TABLE 4.2 RESULTS OF REGRESSION ANALYSIS OF U-WIND COMPONENT IN SUMMER -EXCLUDING THE 1972-
1984 RECORDS, THE SIGNIFICANT REGRESSION VALUES ARE BOLDED.
TABLE 4.3 REGRESSION ANALYSIS OF V-WIND COMPONENT IN SUMMER – EXCLUDING THE 1972-1984
RECORDS
TABLE 4.4 RESULT OF REGRESSION ANALYSIS OF U-WIND COMPONENT IN AUTUMN
TABLE 4.5 RESULT OF REGRESSION ANALYSIS OF V-WIND COMPONENT IN AUTUMN
TABLE 4.6 RESULT OF REGRESSION ANALYSIS OF U-WIND COMPONENT IN SPRING – EXCLUDING THE 1972-
1984 AND 1992 RECORDS
TABLE 4.7 RESULT OF REGRESSION ANALYSIS OF V-WIND COMPONENT IN SPRING –EXCLUDING THE 1972-
1984 AND 1992 RECORDS
TABLE 4.8 THE TIME OF OBSERVED SIGNIFICANT INCREASE OF THE COMPONENT OF AFTERNOON AND
EVENING WINDS
TABLE 4.9 THE CORRELATION BETWEEN EACH AFTERNOON WIND COMPONENT AND THE NEXT READING 83
TABLE 4.10 CORRELATION COEFFICIENT BETWEEN MONTHLY AVERAGED U AND V COMPONENTS OF WIND
AND MAXIMUM TEMPERATURE FOR SEA BREEZE AND NON-SEA BREEZE DAYS
TABLE 5.1 PHYSICAL PARAMETERS OF DIFFERENT LAND COVERS OF METROPOLITAN AREA. 100
TABLE 5.2 STATISTICAL EVALUATION OF SIMULATION OF TEMPERATURE, WIND SPEED AND DIRECTION AT
DIFFERENT LOCATIONS

Abstract

Nearshore processes along the sandy beaches of Adelaide are driven by the prevailing wind and waves. While the narrow entrance of Gulf St. Vincent and the shallow waters attenuate the ocean swell waves, the locally generated southerly to south-westerly waves are behind the net northward littoral drift transport. An important factor in the generation of local waves are the sea breezes, which for Gulf St Vincent result from a combination of a southerly ocean breeze and westerly gulf breezes, and a key question in this time when the climate is said to be changing is whether there is evidence that these sea breezes are changing.

This study, therefore, investigates the existence of long-term changes to the gulf breeze; hereafter referred to as the sea breeze, over the period of August 1955 to June 2008.

In the study of the local climate a set of criteria were developed to define and identify the sea breeze days on which the locally generated coastal winds are generally dominant in the afternoon. Considering the limitation of meteorological observations, the criteria employed the three-hourly near surface data, the 12-hourly upper air levels recorded data, and the surface temperatures of Gulf St. Vincent, provided by Advanced High Resolution Radiometer (AVHRR).

Applying the methods, the period of study is divided into sea breeze and non-sea breeze days, where the characteristic afternoon wind in both categories is analysed. Although the annual percentage of observed sea breeze cases does not show any significant change over the period of the study, the results have demonstrated the presence of an increasing trend in the intensity of afternoon winds, more evidently for the selected sea breeze days.

Through regression analysis of the results, the rise of the southerly component of the wind has been found to have a strong correlation with the surface temperature of the land, whereas the growth of the westerly component was not correlated with any local climate drivers.

Following this important result, the study then went on to determine what might be driving this change. As the importance of urbanization on the climate of wind has been extensively studied by previous researchers, the growth of the Adelaide metropolitan area was conjectured to affect the wind climate at the planetary boundary layer.

A next-generation mesoscale numerical model, Weather Research and Forecasting (WRF), was employed to simulate the climate of the area with and without the metropolitan area of Adelaide, where the city was replaced by the native vegetation of the land. From the simulations it appears that the westerly components of the winds are strongly affected by changes to the nature of the land, due to a combination of changes to the surface roughness length and modification of the surface heat budget components.

The main findings of the statistical and numerical study of the wind climate of Adelaide are:

- 1. The wind climate in and around Gulf St. Vincent has shown a statistically significant change over the last 50 years.
- 2. While there has been no significant change in the number of sea breeze days, the current wind climate has significantly higher wind speeds more evidently on sea breeze days. This is likely to have an important effect on the coast, particularly if the trend continues.
- 3. Through the component-wise analysis of wind, the change in the intensity of south-north component of wind intensity was found to be correlated to the increasing trend of land surface temperature. This is likely to explain one of the key drivers of the change in wind climate.

4. A numerical modelling exercise demonstrated the importance of the growth of the metropolitan area of Adelaide with the change in surface roughness and the change to the surface energy budget being two key elements of the change.

In the end, there is still a need for future study to examine the possible effects of prolonged changes of wind characteristics on the dynamics of the shoreline, particularly in regard to the littoral sediment transport system.

Statement of Originality

I, Zahra Pazandeh Masouleh certify that this work contains no material which has been accepted for the award of any other degree or diploma in my name, in any university or other tertiary institution and, to the best of my knowledge and belief, contains no material previously published or written by another person, except where due reference has been made in the text. In addition, I certify that no part of this work will, in the future, be used in a submission in my name, for any other degree or diploma in any university or other tertiary institution without the prior approval of the University of Adelaide and where applicable, any partner institution responsible for the joint-award of this degree.

I give consent to this copy of my thesis, when deposited in the University Library, being made available for loan and photocopying, subject to the provisions of the Copyright Act 1968.

I also give permission for the digital version of my thesis to be made available on the web, via the University's digital research repository, the Library Search and also through web search engines, unless permission has been granted by the University to restrict access for a period of time.

Signature:

Date:

Acknowledgement

I would like to express my deep and sincere gratitude to my supervisors: Professor David Walker and Professor John Crowther, for their valuable guidance, consistent encouragement, understanding, patience, and most importantly, their friendship throughout the research work.

I am extremely grateful to my main supervisor Prof. David Walker, for his guidance and all the useful discussions and brainstorming sessions, especially during the difficult conceptual development stage. His deep insights helped me at various stages of my research. I also remain indebted for his understanding and support during my maternity leave. My sincere gratitude is reserved for Prof. John Crowther for his invaluable insights and suggestions. I really appreciate his willingness to meet me at short notice every time and going through several drafts of my thesis. I remain amazed that despite his busy schedule, he was able to go through the final draft of my thesis and meet me in less than a week with comments and suggestions on almost every page. He is an inspiration. I also acknowledge the contribution of Dr. Murray Townsend to my thesis and his support as a member of my PhD supervision panel is highly appreciated. I thank Mr. Darren Ray, senior meteorologist at South Australian Regional Climate Services Centre of Bureau of Meteorology for his supports and helps with meteorological data. Sincere thanks are extended to the former computing officer, Dr. Stephen Carr and the school administrative staff, for their help.

I wish to thank my fellow postgraduates in the School of Civil, Environmental and Mining Engineering for their assistance and friendship. Special thanks to my very good friends Li Li, Jaya and Nimasha, who have made this journey so much more enjoyable. I owe a lot to my parents, who encouraged and helped me at every stage of my personal and academic life, and longed to see this achievement come true. Lastly, I would like to dedicate this thesis to my precious daughter, Aida and my loving, supportive, encouraging, and patient husband Arash Asadi whose faithful support through this Ph.D. is so appreciated.

List of Abbreviation

AAO	Antarctic Oscillation
ACST	Australian Central Standard Time
AFWA	Air Force Weather Agency
AHD	Australian Height Datum
AOI	Antarctic Oscillation Indices
ARW	Advanced Research WRF
AVHRR	Advanced Very High Resolution Radiometer
CBD	Central Business District
CGA	Synchrotac Cup Anemometer
CSIRO	Commonwealth Scientific and Industrial Research Organisation
CTL	Control Run
DST	Daylight Saving Time
ENSO	El Niño Southern Oscillation
ETA	Eta Model
FAA	Federal Aviation Administration
FSL	Forecast System Laboratory
GCM	General Circulation Models
GDAS	Global Data Assimilation System
GTS	Global Telecommunications System
IOA	Index Of Agreement
IOD	Indian Ocean Dipole
LH	Latent Heat
LW	Long Wave radiation
MB	Mean Biass Error
MM5	PSU/NCAR mesoscale model
MO	Monin–Obukhov similarity theory
NCAR	National Center for Atmospheric Research
NCEP	National Center for Environmental Prediction
NCEP FNL	global reanalysis datasets (final data analysis) from the US National Centers for Environmental Prediction
NESL	parallel language developed at Carnegie Mellon
NMM	Non-hydrostatic Mesoscale Model
NOAA	National Oceanic and Atmospheric Administration
NRMSE	Normalized root mean square error
NTV	Native Vegetation (model)
NVIS	National Vegetation Information System

PBL	Planetary Boundary Layer
pchip	Piecewise Cubic Hermite Interpolating Polynomial
PTA	Dines Pressure Tube Anemometer
Qv	mixing ratio for water vapour
RAAF	Royal Australian Air Force
RCM	Regional Climate Models
RMSD	Root Mean Square Difference
RMSE	Root Mean Square Error
RSM	Reynolds Stress equation Model
SH	Sensible Heat
SLP	Sea-Level Pressure
SOI	Southern Oscillation Index
SW	Short Wave radiation
UCI	Urban Cool Island
UHI	Urban Heat Island
USGS	U.S. Geological Survey
WMP	Warmer Months Prediction
WPS	WRF pre-processing system
WRF	Weather Research and Forecasting
WSM3	WRF Single Moment 3 classes